

HETEROCYCLES, Vol. 94, No. 8, 2017, pp. 1479 - 1505. © 2017 The Japan Institute of Heterocyclic Chemistry
Received, 27th April, 2017, Accepted, 8th June, 2017, Published online, 14th June, 2017
DOI: 10.3987/COM-17-13736

REACTION OF ELECTRON-DEFICIENT 3-ACETYL-1-ARYLPENT-2-ENE-1,4-DIONES AS A BUILDING BLOCK OF HETEROCYCLES[†]

Shun Kawabata,^a Akihiro Oishi,^a and Hiroshi Nishino^{b*}

^a Department of Chemistry, Graduate School of Science and Technology, Kumamoto University, Kurokami 2-39-1, Chûou-Ku, Kumamoto 860-8555, Japan

^b Department of Chemistry, Faculty of Science, Kumamoto University, Kurokami 2-39-1, Chûou-Ku, Kumamoto 860-8555, Japan

Fax: +81-96-342-3374; E-mail: nishino@kumamoto-u.ac.jp


[†]Dedicated to Dr. Kazu Kurosawa, Professor Emeritus of Kumamoto University, on his 80th birthday.

Abstract – The BF₃-assisted reaction of 3-acetyl-1-arylpent-2-ene-1,4-diones **1a-c** with cyclohexanones and piperidin-4-ones **2a-i** gave unique 3a,6a-dihydrospirofuro[2,3-*d*][1,3]dioxoles **3** in good to high yields. A similar reaction with the 2,3-dihydroquinolin-4(*1H*)-ones did not occur, but the reaction with 4-hydroxychromenone **5** mainly produced 3-furfuryl-4-hydroxychromenone **6** along with furochromenone **7**. The reaction of the electron-deficient pentenedione **1a** as a Michael acceptor with indole, pyrrole, furan, and *N*-methylaniline produced the corresponding 1,4-adducts. Especially, the indole adduct was easily converted by the Paal-Knorr synthesis into the corresponding furanyl-, pyrrolyl-, and thiophenyl-substituted indoles. The reaction details and the structure determination of the products are described.

INTRODUCTION

The synthesis of novel heterocyclic compounds is always a hot topic in organic chemistry because it sometimes can provide enormous opportunities for the discovery of new pharmaceuticals and materials.¹ The cycloaddition reaction is one of the direct and powerful tools for the construction of complex heterocyclic compounds from structurally simple and readily available electron-deficient alkenes as the starting material in a one-step process, and many examples of both ionic and radical reactions have been

found in the literature.² In connection with our previous study, we reported the selective transformation of the endoperoxide intermediate³ obtained by the photosensitized oxygenation of furan **I**⁴ into 3-acetyl-1-phenylpent-2-ene-1,4-dione (**1a**) and oxirane **II** depending on the reaction conditions (Scheme 1).⁵ The electron-deficient pentenedione **1a** is very attractive as a building block in many reactions.⁶ For example, the BF₃-catalyzed reaction of **1a** in the presence of acetylacetone (Hacac) in dry tetrahydrofuran (THF) at rt gave the polyfunctionalized furans **III** and **IV**.^{7a} When the reaction in the absence of Hacac was carried out in wet THF at reflux temperature, bis(furyl)methane **V** was preferentially produced.^{7b} A typical Diels-Alder reaction of **1a** with cyclopentadiene gave a mixture of the *endo/exo* **VI** and bicyclic compound **VII**.^{7b} In order to determine the potential of the pentenedione **1a** as a building block of heterocyclic compounds, we examined the BF₃-assisted reaction using cyclic ketones such as cyclohexanones and piperidin-4-ones **2a-i**. Surprisingly, the reaction did not produce the furyl-substituted products, but the 3a,6a-dihydrospirofuro[2,3-*d*][1,3]dioxoles **3**. We now report the results of the unique cyclization reaction and also describe the BF₃-assisted reaction with 4-hydroxychromenone **5**. In addition, we describe the Michael addition of **1a** with pyrrole, furan, and indole as an electron donor, and also the application using the Paal-Knorr strategy.




Scheme 1. Some Reactions Using the Pentenedione **1a**

RESULTS AND DISCUSSION

Reaction with Cyclohexanones and Piperidin-4-ones

The pentenedione **1a** was easily prepared by the Knoevenagel condensation of phenylglyoxal with Hacac and was not stable in air.^{7b} When the reaction of **1a** with cyclohexanone (3 equiv.) in the presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (1 equiv.) was carried out in dry THF at room temperature, the spirodioxole **3aa** was obtained in 36% yield together with tetrahydrobenzofuran **4aa** (14% yield) (Scheme 2 and Table 1, Entry 1). A prolonged reaction time led to a slight increase in **3aa** (48% yield) (Entry 2). The use of 4 equivalents of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ caused the decomposition of **3aa** (Entry 4). The best yield of **3aa** (58% yield) was realized by the use of **1a** (5 equiv.) and $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (5 equiv.) at 0 °C for 24 h (Entry 3). A similar reaction with 4-*tert*-butylcyclohexanone (**2b**) was conducted at room temperature to afford the corresponding spirodioxole **3ab** as a 47:53 diastereomixture and tetrahydrobenzofuran **4ab** (Entry 5). Interestingly, the reaction at lower than 0 °C led to an increase in the spirodioxole **3ab** even though a longer reaction time was required (Entries 6 and 7). The reaction of the 4-chlorophenyl-substituted pent-2-ene-1,4-dione **1b** with **2b** gave a similar result, but with a much better yield of the corresponding product (Entries 8 and 9). On the other hand, the use of 3-acetyl-1-(4-methoxyphenyl)pent-2-ene-1,4-dione (**1c**) led to an intractable mixture, and tetrahydrobenzofuran **4cb** rather than the spirodioxole **3cb** was produced in 47% yield for the reaction at room temperature (Entry 11). The structures of the products **3** and **4** were characterized by spectroscopic methods and elemental analysis. Fortunately, a single crystal of one of the diastereomers **3bb** was successfully grown from hexane and subjected to an X-ray crystallographic measurement. We then obtained the exact structure of the spirodioxole **3bb** (see experimental section).⁸

Scheme 2. Reaction of Pentenediones **1a-c** with Cyclohexanones **2a,b**Table 1. Reaction of Pentenediones **1a-c** with Cyclohexanes **2a,b** in the Presence of BF_3^a

Entry	1/Ar	2/R	1:2:BF ₃ ^b	Temp/°C	Time/h	Product Yield/% ^c	1 rec./% ^d
1	1a: Ph	2a: H	1:3:1	rt	35	3aa (36) 4aa (14)	-
2	1a	2a	1:3:1	rt	45	3aa (48) 4aa (9)	-

3	1a	2a	1:5:2	0	24	3aa (58)	4aa (9)	-
4	1a	2a	1:5:4	0	17	-	4aa (10)	-
5	1a	2b : <i>t</i> -Bu	1:5:1	rt	8	3ab (36) (dr 47:53) ^e	4ab (19)	-
6	1a	2b	1:5:1	0	24	3ab (55) (dr 40:60) ^e	4ab (16)	-
7	1a	2b	1:5:2	-20	72	3ab (49) (dr 39:61) ^e	4ab (13)	4
8	1b : 4-ClC ₆ H ₄	2b	1:5:1	0	38	3bb (58) (dr 50:50) ^e	4bb (13)	12
9	1b	2b	1:5:2	0	74	3bb (70) (dr 41:59) ^e	4bb (19)	-
10	1c : 4-MeOC ₆ H ₄	2b	1:5:1	0	185	3cb (8) (dr 38:62) ^e	4cb (10)	41
11	1c	2b	1:5:1	rt	24	3cb (2) (dr 50:50) ^e	4cb (47)	-

^a The reaction of **1** (0.5 mmol) with **2a** was carried out in dry THF (0.5 mL).


^b Molar ratio.

^c Isolated yield based on **1** used.

^d Recovery of **1**.

^e Diastereomer ratio based on the ¹H NMR spectrum.

In order to apply the reaction to azacyclic ketones, such as the piperidin-4-ones, we next examined the reaction of pentenedione **1a** with 1-methylpiperidin-4-one (**2c**) (Scheme 3 and Table 2). Although the reaction was carried out in dry THF at room temperature according to the conditions described above, the reaction did not proceed at all (Entry 1). The reaction at reflux temperature afforded the corresponding azaspirodioxole **3ac** in a poor yield (Entry 2). When the reaction was carried out in dichloromethane (DCM) at room temperature, the yield of **3ac** increased (Entry 3). The reaction was then optimized and a synthetically acceptable yield of **3ac** was achieved at 0 °C under neat conditions (Entry 6). Since the viscosity of **2c** at -20 °C was much higher than that at 0 °C, a small amount of DCM (0.05 mL) was added to the mixture and the reaction was then conducted at -20 °C to give almost the same result (Entry 7).



Scheme 3. Reaction of Pentenedione **1a** with Piperidin-4-ones **2c-i**

Table 2. Reaction of Pentenedione **1a** with Piperidin-4-ones **2c-i** in the Presence of BF₃^a

Entry	2/R	1:2:BF₃^b	Solvent/mL	Temp/°C	Time/h	Product Yield/% ^c
1	2c : Me	1:1:1	THF/5	rt	24	no reaction
2	2c	1:1:3	THF/5	reflux	6	3ac (14) ^d
3	2c	1:1:1	DCM/5	rt	12	3ac (25) ^d
4	2c	1:1:5	DCM/5	rt	6	3ac (61)
5	2c	1:1:5	neat	rt	2.5	3ac (74)
6	2c	1:1:5	neat	0	2.5	3ac (93)
7	2c	1:1:5	DCM/0.05	-20	2.5	3ac (94)
8	2d : Et	1:1:5	neat	0	2.5	3ad (71)
9	2d	1:1:5	DCM/0.05	-40	3	3ad (88)
10	2e : Pr	1:1:5	neat	0	2.5	3ae (68)
11	2e	1:1:5	DCM/0.05	-20	2.5	3ae (77)
12	2f : Bn	1:1:5	neat	0	2.5	3af (68)
13	2f	1:1:5	DCM/0.05	-40	4	3af (89)
14	2g : Ac	1:1:5	neat	0	2.5	3ag (79)
15	2g	1:1:5	DCM/0.05	-20	2.5	3ag (quant)
16	2h : Bz	1:1:5	neat	0	2.5	3ah (85)
17	2h	1:1:5	DCM/0.05	-20	3	3ah (94)
18	2i : Cbz	1:1:5	neat	0	3	3ai (94)

^a The reaction of **1a** (0.5 mmol) with **2** was carried out in dry solvent or under neat conditions.

^b Molar ratio.

^c Isolated yield based on **1a** used.

^d A small amount of homocyclization products **I** and **V** was isolated.


Having succeeded in the cycloaddition of the pentenedione **1a** with the azacyclohexanone **2c**, we turned our attention to the reaction with other piperidin-4-ones (Table 2, Entries 8-18). As a result, both the *N*-alkyl- **2d-f** and *N*-acyl-substituted piperidin-4-ones **2g-i** were tolerated during the cycloaddition reaction under neat or low temperature conditions, producing the corresponding azaspirodioxoles **3ad-3ai** in high yields. A similar reaction with the 1-methyl- and 1-tosyl-2,3-dihydroquinolin-4(1*H*)-ones did not occur, but the homocyclization product **V** of **1a** was only produced.

The NMR spectrum of *N*-acetylspirodioxole **3ag** in CDCl₃ at room temperature deserves comments. In the ¹H NMR spectrum, two signals appeared at δ 2.13 (s) and 2.11 (s) assigned to the *N*-acetyl group and most of the signals in the ¹³C NMR spectrum each showed two peaks. When the NMR spectrum was measured in DMSO-*d*₆ at 80 °C, the *N*-acetyl group in the ¹H NMR spectrum and the signals that appeared for each of the two peaks in the ¹³C NMR spectrum collapsed into one peak due to existence of the conformational isomer in CDCl₃. Although a similar phenomenon was not observed in other azaspirodioxoles, the methylene peaks of the *N*-acylspirodioxoles **3ah** and **3ai** in the ¹³C NMR spectrum

appeared as a weak signal on the NMR time scale, probably depending on the flipping velocity of the azacyclohexane ring.

Reaction with 4-Hydroxy-2*H*-chromen-2-one

With the unique spirodioxole formation in hand, the reaction using 4-hydroxy-2*H*-chromen-2-one (**5**) was next investigated. When the reaction was carried out at the molar ratio of **1a**:**5**:BF₃ = 1:3:1 in DCM (1.0 mL) at rt, unfortunately, the corresponding dioxole was not isolated, but furfurylchromenone **6** and furochromenone **7** were obtained together with bis(furyl)methane **V** as the homocyclization product of **1a** (Scheme 4 and Table 3, Entry 1).^{7b} In order to prevent the production of **V**, the pentenedione **1a** was dropwise added to the mixture of **5** and BF₃ at reflux temperature. As a result, the yield of **6** was improved (Entry 2), but bis(furyl)methane **V** was still produced. We then scrutinized the solvent and reaction temperature, and eventually, the use of chlorobenzene at 130 °C led to the maximum yield of **6** and suppressed the production of **V** (Entry 6).



Scheme 4. Reaction of Pentenedione **1a** with 4-Hydroxy-2*H*-chromene-2-one (**5**)

Table 3. Reaction of Pentenedione **1a** with 4-Hydroxychromenone (**5**) in the Presence of BF₃^a

Entry	Solvent/mL	Temperature/°C	Time/min	Product Yield/% ^b		
				6	7	V ^c
1	DCM/1	rt	70	24	14	16
2 ^d	DCM/1	40	50	41	1	12
3 ^d	DCE/1	80	20	51	17	3
4 ^d	C ₆ H ₅ Cl/1	100	10	57	12	14
5 ^d	C ₆ H ₅ Cl/1	130	5	69	13	7
6 ^d	C ₆ H ₅ Cl/0.5	130	3	77	13	5

^a The reaction of **1a** (0.3 mmol) with **5** (0.9 mmol) was carried out in the presence of BF₃·OEt₂ (0.3 mmol).


^b Isolated yield based on **1a** used.

^c Bis(3-acetyl-5-phenyl-2-furyl)methane (**V**) as a homocyclization product of **1a**.


^d Pentenedione **1a** (0.3 mmol) in solvent (0.3 mL) was dropwise added to the mixture of **5** (0.9 mmol) and BF₃·OEt₂ (0.3 mmol) in solvent.

Proposed Mechanism for the Formation of the Products

The BF_3 -assisted cyclization of pentenediones **1** with cyclic ketones **2** could be similarly understood by the previously reported literature as depicted in Scheme 5.⁷ The production of two diastereomers in the reaction using 4-*tert*-butylcyclohexanone (**2b**) would be caused by exclusive occupation of the *tert*-butyl group at the equatorial position. That is, the carbonyl group of the cyclohexanone equatorial fixed *tert*-butyl group might be attacked by the pentenedione **1** from the equatorial or axial direction, affording the corresponding diastereomeric dioxoles. In the case of **1c** (Ar = 4-MeOC₆H₄) (Table 1, Entries 10 and 11), the production of the spirodioxole **3cb** was suppressed because the electron-donating methoxy group might change the electrostatic potential of **1c**.




Scheme 5. Proposed Mechanism for the Formation of Dioxoles **3**



Scheme 6. Proposed Mechanism for the Formation of Tetrahydrobenzofurans **4**

On the other hand, the production of the by-products **4** would be explained by an aldol-type condensation briefly shown in Scheme 6. This reaction could not be retarded under all of the conditions. However, the corresponding by-product, such as tetrahydrobenzofuran **4**, was not produced during the reaction using the azacyclohexanones **2c-i**.


The furfurylchromenone **6** would be formed by the reaction of methylenedihydrofuranol **A** prepared by the enolization of **1a** followed by in situ cyclization (Scheme 7).⁷ It was considered that the by-product furochromenone **7** would be obtained by a similar reaction sequence as shown in Scheme 6.



Scheme 7. Plausible Mechanism for the Formation of Furfurylchromenone **6**

Michael Addition Using Indole, Pyrrole, Furan, and *N*-Methylaniline

In order to further study the reactivity of the electron-deficient pentenedione **1a** as a Michael acceptor, we next investigated the reaction with indole as an electron-donor (Scheme 8).⁹ The reaction was examined in various solvents and the Michael adduct **8** was obtained in all the solvents (see Experimental section). The best yield of **8** (97% yield) was realized in acetonitrile at reflux temperature for 9.5 h. A similar reaction with pyrrole gave the 1,4-adduct **9** in 84% yield.




Scheme 8. Michael Addition of **1a** with Indole, Pyrrole, Furan, and *N*-Methylaniline

In addition, the use of a one-half equivalent of pyrrole provided the 2,5-disubstituted adduct in 79% yield (see Experimental section). On the other hand, the reaction with furan did not proceed in acetonitrile, but the corresponding adduct **10** (76% yield) was obtained in acetic acid at room temperature. *N*-Methylaniline also afforded the 1,4-adduct **11** (75% yield) in boiling acetic acid.

Conversion into Heterocycle-substituted Indole Derivatives via Paal-Knorr Strategy

With the 1,4-dicarbonyl-functionalized indole **8** in hand, we finally embarked on the synthesis of the five-membered heterocycle-substituted indoles using the Paal-Knorr strategy (Scheme 9).¹⁰ Since it was reported that the treatment of **1a** with indole in the presence of methanesulfonic acid gave the 3-furyl-substituted indole,^{10c} we initially examined the BF₃-catalyzed reaction of **1a** with indole in DCM at reflux temperature. Pleasingly, the reaction finished in only 30 min¹¹ and the reported product **12** (88% yield) was obtained during the formation of **8**. The Michael adduct **8** underwent the typical Paal-Knorr reaction with ammonium acetate to produce the desired pyrrole-substituted indole **13a** in quantitative yield without purification. Use of benzylamine and ethanolamine also gave the desired products **13b** and **13c** in high yields. Similarly, the treatment of **8** with Lawesson's reagent in toluene at 50 °C afforded the thienyl-substituted indole **14** (49% yield) together with **12** (23% yield).¹²



Scheme 9. Synthesis of Heterocycle-substituted Indole Derivatives

CONCLUSION

We have accomplished the unique synthesis of spirodioxoles and azaspirodioxoles **3** by the cycloaddition of pentenediones **1** with cyclohexanones and piperidin-4-ones **2**. The reaction with chromenone **5** did not

give the spirodioxole, but furfurylchromenone **6** along with furochromenone **7**. There is no precedent for the synthesis of bicyclo-spirodioxoles **3** except for a similar protecting diol of sugars¹³ and kendomycin analogues which have a cytotoxic property.¹⁴ We also demonstrated the usefulness of pentenedione **1a** as a convenient building block in the Michael addition followed by the Paal-Knorr heterocycle synthesis using indole.


EXPERIMENTAL

Measurements. Melting points were taken using a Yanagimoto micromelting point apparatus and are uncorrected. The NMR spectra were recorded using a JNM ECX 500 or AL300 FT-NMR spectrometer at 500 MHz for the ¹H and at 125 MHz for ¹³C, with tetramethylsilane as the internal standard. The chemical shifts are reported as δ values (ppm) and the coupling constants in Hz. The following abbreviations are used for the multiplicities: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; and brs, broad singlet for the ¹H NMR spectrum. The IR spectra were measured in CHCl₃ or KBr using a Shimadzu 8400 FT IR spectrometer and expressed in cm⁻¹. The high-resolution mass spectra and the elemental analyses were performed at the Instrumental Analysis Center, Kumamoto University, Kumamoto, Japan. The X-ray analysis was performed by a Rigaku RAXIS RAPID imaging plate area detector with graphite monochromated Mo-K α radiation.


Materials. The 3-acetyl-1-aryl-2-pentene-1,4-diones **1a-c** and piperidin-4-ones **2e**, **2h**, **2i** were prepared according to the literature method.¹⁴ Cyclohexanone (**2a**), 4-*tert*-butylcyclohexanone (**2b**), 1-methylpiperidin-4-one (**2c**), 1-ethylpiperidin-4-one (**2d**), 1-benzylpiperidin-4-one (**2f**), 4-hydroxy-2H-chromen-4-one (**5**), indole, pyrrole, furan, and *N*-methylaniline were purchased from Tokyo Kasei Co., Ltd., and 1-acetylpiperidin-4-one (**2g**) and the boron trifluoride diethyl ether complex were from Wako Pure Chemical Ind., Ltd., and used as received. Flash column chromatography was performed on silica gel 60N (40-50 mm), which was purchased from Kanto Chemical Co., Inc., and preparative thin layer chromatography (TLC) on Wakogel B-5F from Wako Pure Chemical Ind., Ltd. The solvents were commercially available first grade and used as received.

Reaction of 1 with Cyclohexanones 2a and 2b in the Presence of BF₃·OEt₂. A solution of pentenedione **1a** (216.8 mg, 1.0 mmol) and cyclohexanone (**2a**) (490.6 mg, 2.5 mmol) in dry THF (1.0 mL) was cooled at 0 °C under argon, and BF₃·Et₂O (254 μ L, 2.0 mmol) was slowly added using a syringe. After the reaction was completed, water (10 mL), brine (10 mL), and a saturated aqueous solution of NaHCO₃ (30 mL) were then added, and the aqueous mixture was extracted with CHCl₃ (3 \times 20 mL). The combined extracts were washed with water (20 mL), dried over anhydrous sodium sulfate, and then concentrated to dryness. The residue was purified by column chromatography on silica gel eluting with


EtOAc/hexane (1:5 v/v), giving the desired spirodioxole **3aa** (183 mg, 58%) and tetrahydrobenzofuran **4aa** (25.6 mg, 9%) (Table 1, Entry 3).



6'-Acetyl-5'-methyl-3a'-phenyl-3a',6a'-dihydrospiro[cyclohexane-1,2'-furo[2,3-*d*][1,3]dioxole] (3aa): Reaction time, 24 h; yield 58%; pale yellow oil; $R_f = 0.44$ (EtOAc-hexane 1:4 v/v); IR (CHCl₃) ν 2939 (PhH), 1672 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.46-7.38 (5H, m, arom. H), 5.31 (1H, s, H-6a'), 2.40 (3H, s, Me), 2.29 (3H, s, Ac), 1.86-1.45 (8H, m, 4×CH₂), 1.44 (2H, quin, $J = 4.5$ Hz, CH₂); ¹³C NMR (125 MHz, CDCl₃) δ 194.7 (C=O), 171.1 (C-5'), 137.6 (arom. C), 129.2 (arom. CH), 128.5 (2C) (arom. CH), 125.0 (2C) (arom. CH), 114.9 (C-3a'), 114.8 (C-6'), 114.2 (C-2'), 88.7 (C-6a'), 37.2 (2×CH₂), 29.4 (O=CMe), 24.8, (2×CH₂), 23.9 (CH₂), 15.3 (Me); FAB HRMS (acetone/NBA): calcd for C₁₉H₂₂O₄Na 337.1416 (M+Na); Found 337.1395.




6'-Acetyl-4-(*tert*-butyl)-5'-methyl-3a'-phenyl-3a',6a'-dihydrospiro[cyclohexane-1,2'-furo[2,3-*d*][1,3]dioxole] (3ab): Reaction time, 24 h; yield 22%; colorless needles (from hexane); mp 84-87 °C; $R_f = 0.60$ (EtOAc-hexane 1:4 v/v); IR (KBr) ν 2957 (PhH), 1672 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.46-7.39 (5H, m, arom. H), 5.32 (1H, s, H-6a'), 2.41 (3H, s, Me), 2.30 (3H, s, Ac), 1.96-1.83 (4H, m, 2×CH₂), 1.73-1.64 (2H, m, CH₂), 1.40 (1H, dq, $J = 10.0, 3.0$ Hz, H-CH), 1.30-1.25 (1H, m, H-CH), 1.06 (1H, tt, $J = 12.5, 3.0$ Hz, H-4), 0.87 (3H×3, s, 3×Me); ¹³C NMR (125 MHz, CDCl₃) δ 194.9 (C=O), 171.2 (C-5'), 137.6 (arom. C), 129.3 (arom. CH), 128.6 (2C) (arom. CH), 125.1 (2C) (arom. CH), 115.0 (C-3a'), 114.7 (C-6'), 113.9 (C-2'), 88.9 (C-6a'), 46.7 (C-4), 37.2 (CH₂), 36.9 (CH₂), 32.3 (C(Me)₃), 29.5 (O=CMe), 27.6 (3×Me), 24.8 (CH₂), 24.7 (CH₂), 15.3 (Me); FAB HRMS (acetone/NBA): calcd for C₂₃H₃₀O₄Na 393.2042 (M+Na); Found 393.2014.




6'-Acetyl-4-(*tert*-butyl)-5'-methyl-3a'-phenyl-3a',6a'-dihydrospiro[cyclohexane-1,2'-furo[2,3-*d*][1,3]dioxole] (3ab'): Reaction time, 24 h; yield 33%; colorless oil; $R_f = 0.54$ (EtOAc-hexane 1:4 v/v); IR (CHCl₃) ν 2955 (PhH), 1672 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.46-7.39 (5H, m, arom. H), 5.29 (1H, s, H-6a'), 2.41 (3H, s, Me), 2.29 (3H, s, Ac), 1.97-1.95, 1.90-1.83, 1.78-1.72 (5H, m, CH₂), 1.58 (1H, dt, $J = 13.0, 4.0$ Hz, H-CH), 1.49 (1H, dq, $J = 13.0, 4.0$ Hz, H-CH), 1.40 (1H, dq, $J = 13.0, 4.0$ Hz, H-CH),

1.06 (1H, tt, $J = 13.0, 4.0$ Hz, H-4), 0.87 (3H×3, s, 3×Me); ^{13}C NMR (125 MHz, CDCl_3) δ 194.7 (C=O), 171.2 (C-5'), 137.7 (arom. C), 129.3 (arom. CH), 128.6 (2C) (arom. CH), 125.1 (2C) (arom. CH), 115.1 (C-3a'), 115.0 (C-6'), 114.5 (C-2'), 88.6 (C-6a'), 47.0 (C-4), 37.5 (CH_2), 37.4 (CH_2), 32.3 ($\text{C}(\text{Me})_3$), 29.4 (O=CMe), 27.6 (3×Me), 24.9 (CH_2), 24.8 (CH_2), 15.3 (Me); FAB HRMS (acetone/NBA): calcd for $\text{C}_{23}\text{H}_{30}\text{O}_4\text{Na}$ 393.2042 (M+Na); Found 393.2029.




6'-Acetyl-4-(tert-butyl)-3a'-(4-chlorophenyl)-5'-methyl-3a',6a'-dihydrospiro[cyclohexane-1,2'-furo[2,3-d][1,3]dioxole] (3bb): Reaction time, 74 h; yield 29%; colorless prisms (from hexane); mp 140-143 °C; $R_f = 0.67$ (EtOAc-hexane 1:4 v/v); IR (KBr) ν 2953 (PhH), 1682 (C=O); ^1H NMR (500 MHz, CDCl_3) δ 7.37 (4H, s, arom. H), 5.28 (1H, s, H-6a'), 2.39 (3H, s, Me), 2.30 (3H, s, Ac), 1.92-1.86 (4H, m, 2× CH_2), 1.73-1.63 (2H, m, CH_2), 1.40 (1H, dq, $J = 12.3, 3.5$ Hz, H-CH), 1.25 (1H, dq, $J = 12.3, 3.5$ Hz, H-CH), 1.06 (1H, tt, $J = 11.5, 3.0$ Hz, H-4), 0.87 (3H×3, s, 3×Me); ^{13}C NMR (125 MHz, CDCl_3) δ 194.7 (C=O), 171.0 (C-5'), 136.2, 135.3 (Cl-C, arom. C), 128.8 (2C) (arom. CH), 126.6 (2C) (arom. CH), 115.2 (C-3a'), 114.1 (C-6'), 113.9 (C-2'), 88.9 (C-6a'), 46.6 (C-4), 37.2 (CH_2), 36.8 (CH_2), 32.2 ($\text{C}(\text{Me})_3$), 29.5 (O=CMe), 27.5 (3×Me), 24.8 (CH_2), 24.6 (CH_2), 15.3 (Me); Anal Calcd for $\text{C}_{23}\text{H}_{29}\text{ClO}_4$: C, 67.89; H, 7.09. Found C, 68.22; H, 7.22. X-Ray crystallographic data: empirical formula $\text{C}_{23}\text{H}_{29}\text{ClO}_4$; formula weight 404.18; colorless prisms; orthorhombic; space group Pbca; cell lengths $a = 11.4756(7)$, $b = 19.2308(12)$, $c = 19.830(1)$ Å; cell volume 4376.2(4) Å³; formula units per cell $Z = 8$; total data collected 5352; $R = 0.0557$; $R_w = 0.1068$; GOF = 3.006.




6'-Acetyl-4-(tert-butyl)-3a'-(4-chlorophenyl)-5'-methyl-3a',6a'-dihydrospiro[cyclohexane-1,2'-

furo[2,3-*d*][1,3]dioxole (3bb'): Reaction time, 74 h; yield 41%; colorless oil; $R_f = 0.56$ (EtOAc-hexane 1:4 v/v); IR (CHCl₃) ν 1672 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.39-7.35 (4H, m, arom. H), 5.26 (1H, s, H-6a'), 2.40 (3H, s, Me), 2.27 (3H, s, Ac), 1.95-1.72 (5H, m, \underline{H} -CH, 2 \times CH₂), 1.58 (1H, dt, $J = 13.0, 3.0$ Hz, \underline{H} -CH), 1.47 (1H, dq, $J = 13.0, 3.0$ Hz, \underline{H} -CH), 1.38 (1H, dq, $J = 13.0, 3.0$ Hz, \underline{H} -CH), 1.06 (1H, tt, $J = 13.0, 3.0$ Hz, H-4), 0.87 (3H \times 3, s, 3 \times Me); ¹³C NMR (125 MHz, CDCl₃) δ 194.5 (C=O), 170.9 (C-5'), 136.3, 135.2 (C-C, arom. C), 128.7 (2C) (arom. CH), 126.6 (2C) (arom. CH), 115.1 (C-3a'), 114.4 (C-2', C-6'), 88.5 (C-6a'), 46.9 (C-4), 37.4 (CH₂), 37.3 (CH₂), 32.3 (C(Me)₃), 29.4 (O=CMe), 27.6 (3 \times Me), 24.9 (CH₂), 24.8 (CH₂), 15.3 (Me); FAB HRMS (acetone/NBA): calcd for C₂₃H₂₉ClO₄Na 427.1652 (M+Na); Found 427.1650.




6'-Acetyl-4-(tert-butyl)-3a'-(4-methoxyphenyl)-5'-methyl-3a',6a'-dihydrospiro[cyclohexane-1,2'-furo[2,3-*d*][1,3]dioxole] (3cb): Reaction time, 185 h; yield 3%; colorless oil; $R_f = 0.45$ (EtOAc-hexane 1:4 v/v); IR (CHCl₃) ν 2961 (PhH), 1709, 1670 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.38-7.36 (2H, m, arom. H), 6.92-6.90 (2H, m, arom. H), 5.28 (1H, s, H-6a'), 3.82 (3H, s, OMe), 2.39 (3H, s, Me), 2.30 (3H, s, Ac), 1.93-1.81 (4H, m, 2 \times CH₂), 1.72-1.63 (2H, m, CH₂), 1.40 (1H, dq, $J = 12.0, 3.0$ Hz, \underline{H} -CH), 1.27-1.24 (1H, m, \underline{H} -CH), 1.06 (1H, tt, $J = 12.0, 3.0$ Hz, H-4), 0.87 (3H \times 3, s, 3 \times Me); ¹³C NMR (125 MHz, CDCl₃) δ 194.9 (C=O), 171.2 (C-5'), 160.3 (MeO-C), 129.8 (arom. C), 126.5 (2C) (arom. CH), 114.8 (C-3a'), 114.7 (C-6'), 113.9 (C-2'), 113.9 (2C) (arom. CH), 88.8 (C-6a'), 55.3 (OMe), 46.7 (C-4), 37.3 (CH₂), 36.9 (CH₂), 32.3 (C(Me)₃), 29.5 (O=CMe), 27.6 (3 \times Me), 24.8 (CH₂), 24.7 (CH₂), 15.3 (Me); FAB HRMS (acetone/NBA): calcd for C₂₄H₃₂O₅Na 423.2174 (M+Na); Found 423.2169.




6'-Acetyl-4-(tert-butyl)-3a'-(4-methoxyphenyl)-5'-methyl-3a',6a'-dihydrospiro[cyclohexane-1,2'-furo[2,3-*d*][1,3]dioxole] (3cb'): Reaction time, 185 h; yield 5%; colorless oil; $R_f = 0.42$ (EtOAc-hexane 1:4 v/v); IR (CHCl₃) ν 2959 (PhH), 1710, 1670 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.38-7.36 (2H, m, arom. H), 6.94-6.91 (2H, m, arom. H), 5.26 (1H, s, H-6a'), 3.82 (3H, s, OMe), 2.39 (3H, s, Me), 2.30 (3H, s, Ac), 1.95-1.82 (5H, m, \underline{H} -CH, 2 \times CH₂), 1.72-1.63 (1H, m, \underline{H} -CH), 1.48 (1H, dq, $J = 12.0, 3.5$ Hz, \underline{H} -CH), 1.40 (1H, dq, $J = 12.0, 3.5$ Hz, \underline{H} -CH), 1.06 (1H, tt, $J = 12.0, 3.5$ Hz, H-4), 0.87 (3H \times 3, s, 3 \times Me); ¹³C NMR (125 MHz, CDCl₃) δ 194.8 (C=O), 171.0 (C-5'), 160.3 (MeO-C), 129.9 (arom. C), 126.5 (2C) (arom. CH), 115.2 (C-3a'), 114.7 (C-6'), 114.5 (C-2'), 113.9 (2C) (arom. CH), 88.5 (C-6a'),


55.4 (OMe), 47.0 (C-4), 37.5 (2×CH₂), 32.3 (C(Me)₃), 29.4 (O=CMe), 27.7 (3×Me), 24.9 (2×CH₂), 15.4 (Me); FAB HRMS (acetone/NBA): calcd for C₂₄H₃₂O₅Na 423.2174 (M+Na); Found 423.2140.



(Z)-4-Hydroxy-3-(3-phenyl-4,5,6,7-tetrahydrobenzofuran-2-yl)pent-3-en-2-one (4aa): Reaction time, 24 h; yield 9%; colorless cubes (from methanol); mp 92-97 °C; *R_f* = 0.75 (Et₂O-hexane 1:1 v/v); IR (KBr) ν 2941 (PhH), 1605 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 16.81 (1H, s, OH), 7.35-7.32 (2H, m, arom. H), 7.25-7.21 (3H, m, arom. H), 2.65 (2H, t, *J* = 5.5 Hz, CH₂), 2.51 (2H, t, *J* = 5.5 Hz, CH₂), 1.91 (6H, s, 2×Me), 1.91-1.89 (2H, m, CH₂), 1.79-1.76 (2H, m, CH₂); ¹³C NMR (125 MHz, CDCl₃) δ 193.7 (C=O), 151.0 (C-7a), 143.1 (C-2), 133.5 (arom. C), 128.6 (2C) (arom. CH), 127.8 (arom. CH), 126.6 (2C) (arom. CH), 124.8 (C-3), 117.1 (C-3a), 105.6 (C-3'), 23.9 (2×Me), 23.3 (CH₂), 23.2 (CH₂), 22.8 (CH₂), 22.1 (CH₂); Anal Calcd for C₁₉H₂₀O₃: C, 77.00; H, 6.80. Found C, 76.85; H, 6.89.




(Z)-3-(5-(tert-Butyl)-3-phenyl-4,5,6,7-tetrahydrobenzofuran-2-yl)-4-hydroxypent-3-en-2-one (4ab): Reaction time, 24 h; yield 16%; colorless oil; *R_f* = 0.78 (EtOAc-hexane 1:4 v/v); IR (CHCl₃) ν 2963 (PhH), 1605 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 16.80 (1H, s, OH), 7.37-7.21 (5H, m, arom. H), 2.74 (1H, dd, *J* = 16.5, 4.5 Hz, H-CH), 2.61 (1H, m, H-CH), 2.49 (1H, m, H-CH), 2.31 (1H, t, *J* = 13.5, Hz, H-5), 2.11 (1H, m, H-CH), 2.05 (3H, s, Me), 1.76 (3H, s, Me), 1.55-1.43 (2H, m, CH₂), 0.96 (9H, s, 3×Me); ¹³C NMR (125 MHz, CDCl₃) δ 194.5, 192.9 (C=O), 151.0 (C-7a), 143.4 (C-2), 133.5 (arom. C), 128.6 (2C) (arom. CH), 127.9 (arom. CH), 126.6 (2C) (arom. CH), 125.1 (C-3), 117.3 (C-3a), 105.6 (C-3'), 45.3 (C-5), 32.5 (C(Me)₃), 27.5 (3×Me), 24.3 (CH₂), 24.2 (CH₂), 24.0 (CH₂), 23.3 (2×Me); FAB HRMS (acetone/NBA): calcd for C₂₃H₂₈O₃ 352.2038 (M); Found 352.2052.



(Z)-3-(5-(tert-Butyl)-3-(4-chlorophenyl)-4,5,6,7-tetrahydrobenzofuran-2-yl)-4-hydroxypent-3-en-2-one (4bb): Reaction time, 74 h; yield 19%; colorless prisms (from CH₂Cl₂); mp 113-116 °C; *R_f* = 0.70 (EtOAc-hexane 1:4 v/v); IR (CHCl₃) ν 2963 (PhH), 1607 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 16.82 (1H, s, OH), 7.33-7.32 (2H, m, arom. H), 7.17-7.15 (2H, m, arom. H), 2.73 (1H, dd, *J* = 16.0, 4.5 Hz, H-CH), 2.60 (1H, t, *J* = 15.0 Hz, H-CH), 2.45 (1H, d, *J* = 13.5 Hz, H-CH), 2.28 (1H, t, *J* = 12.5, Hz, H-5), 2.11 (1H, m, H-CH), 2.05 (3H, s, Me), 1.76 (3H, s, Me), 1.52-1.46 (2H, m, CH₂), 0.96 (9H, s,

3×Me); ^{13}C NMR (125 MHz, CDCl_3) δ 194.5, 192.8 (C=O), 151.3 (C-7a), 143.6 (C-2), 132.5, 132.0 (Cl-C, arom. C), 129.1 (2C) (arom. CH), 128.9 (2C) (arom. CH), 124.0 (C-3), 117.1 (C-3a), 105.3 (C-3'), 45.3 (C-5), 32.6 (C(Me) $_3$), 27.5 (3×Me), 24.2 (CH $_2$), 24.0 (CH $_2$), 23.5 (CH $_2$), 23.2 (2×Me); FAB HRMS (acetone/NBA): calcd for $\text{C}_{23}\text{H}_{27}\text{ClO}_3$ 386.1649 (M); Found 386.1646.




(Z)-3-(5-(tert-Butyl)-3-(4-methoxyphenyl)-4,5,6,7-tetrahydrobenzofuran-2-yl)-4-hydroxypent-3-en-2-one (4cb): Reaction time, 24 h; yield 47%; pale yellow oil; R_f = 0.62 (EtOAc-hexane 1:4 v/v); IR (CHCl_3) ν 2961 (PhH), 1705, 1612 (C=C–C=O); ^1H NMR (500 MHz, CDCl_3) δ 16.79 (1H, s, OH), 7.16-7.14 (2H, m, arom. H), 6.90-6.89 (2H, m, arom. H), 3.81 (3H, s, OMe), 2.72 (1H, dd, J = 16.0, 5.0 Hz, $\underline{\text{H}}$ -CH), 2.60 (1H, t, J = 11.5 Hz, $\underline{\text{H}}$ -CH), 2.47 (1H, d, J = 13.5 Hz, $\underline{\text{H}}$ -CH), 2.28 (1H, t, J = 12.5, Hz, H-5), 2.09 (1H, dd, J = 11.5, 4.0 Hz, $\underline{\text{H}}$ -CH), 2.04 (3H, s, Me), 1.77 (3H, s, Me), 1.52-1.45 (2H, m, CH $_2$), 0.95 (9H, s, 3×Me); ^{13}C NMR (125 MHz, CDCl_3) δ 194.4, 193.0 (C=O), 158.3 (OMe-C), 150.9 (C-7a), 142.9 (C-2), 129.0 (2C) (arom. CH), 125.9, 124.6 (C-3, arom. C), 117.3 (C-3a), 114.1 (2C) (arom. CH), 105.7 (C-3'), 55.2 (OMe), 45.3 (C-5), 32.5 (C(Me) $_3$), 27.5 (3×Me), 24.3 (CH $_2$), 24.2 (CH $_2$), 24.0 (CH $_2$), 23.3 (2×Me); FAB HRMS (acetone/NBA): calcd for $\text{C}_{24}\text{H}_{30}\text{O}_4$ 382.2144 (M); Found 382.2154.

Reaction of 1a with Piperidin-4-ones 2c-i in the Presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$. A solution of pentenedione **1a** (106.7 mg, 0.5 mmol) and 1-methylpiperidin-4-one (**2c**) (57.6 mg, 0.5 mmol) was cooled at 0 °C under argon, and $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (315 mL, 2.5 mmol) was slowly added using a syringe. After the reaction was completed, a saturated aqueous solution of NaHCO_3 (30 mL) was then added and the aqueous mixture was extracted with CHCl_3 (3×20 mL). The combined extracts were dried over anhydrous sodium sulfate, then concentrated to dryness. The residue was purified by column chromatography on silica gel eluting with $\text{CHCl}_3/\text{MeOH}$ (95:5 v/v), giving the desired azaspirodioxole **3ac** (149.6 mg, 93%) (Table 2, Entry 6).




6-Acetyl-1',5-dimethyl-3a-phenyl-3a,6a-dihydrospiro[furo[2,3-d][1,3]dioxole-2,4'-piperidine] (3ac): Reaction time, 2.5 h; yield 93%; yellow oil; R_f = 0.22 (CHCl_3 -MeOH 95:5 v/v); IR (CHCl_3) ν 1674 (C=C–C=O); ^1H NMR (500 MHz, CDCl_3) δ 7.45-7.39 (5H, m, arom. H), 5.34 (1H, s, H-6a), 2.64-2.47 (4H, m, 2×NCH $_2$), 2.40 (3H, s, Me), 2.31 (3H, s, NMe), 2.29 (3H, s, Ac), 2.03-1.86 (4H, m, 2×CH $_2$); ^{13}C NMR (125 MHz, CDCl_3) δ 194.4 (C=O), 171.0 (C-5), 137.1 (arom. C), 129.2, 128.5 (2C) (arom. CH),

124.9 (arom. CH), 114.6 (C-3a), 113.9 (C-6), 112.3 (C-2), 88.7 (C-6a), 53.2 (2×NCH₂), 45.6 (NMe), 36.7, 36.5 (2×CH₂), 29.3 (Ac), 15.1 (Me); FAB HRMS (acetone/NBA): calcd for C₁₉H₂₄O₄N 330.1705 (M+H); Found 330.1712.




6-Acetyl-1'-ethyl-5-methyl-3a-phenyl-3a,6a-dihydrospiro[furo[2,3-*d*][1,3]dioxole-2,4'-piperidine]

(3ad): Reaction time, 3.0 h; yield 88%; yellow oil; $R_f = 0.27$ (CHCl₃-MeOH 95:5 v/v); IR (CHCl₃) ν 1674 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.46-7.39 (5H, m, arom. H), 5.34 (1H, s, H-6a), 2.68-2.59 (4H, m, 2×NCH₂), 2.46 (2H, q, $J = 7.3$ Hz, NCH₂CH₃), 2.40 (3H, s, Me), 2.30 (3H, s, Ac), 2.03-1.87 (4H, m, 2×CH₂), 1.10 (3H, t, $J = 7.3$ Hz, NCH₂CH₃); ¹³C NMR (125 MHz, CDCl₃) δ 194.4 (C=O), 170.9 (C-5), 137.2 (arom. C), 129.2, 128.4 (2C) (arom. CH), 124.9 (arom. CH), 114.6 (C-3a), 113.9 (C-6), 112.8 (C-2), 88.6 (C-6a), 51.6, 50.6 (2×NCH₂, NCH₂CH₃), 36.7, 36.5 (2×CH₂), 29.3 (Ac), 15.1 (Me), 12.2 (NCH₂CH₃); FAB HRMS (acetone/NBA): calcd for C₂₀H₂₆O₄N 344.1862 (M+H); Found 344.1867.



6-Acetyl-5-methyl-3a-phenyl-1'-propyl-3a,6a-dihydrospiro[furo[2,3-*d*][1,3]dioxole-2,4'-piperidine]


(3ae): Reaction time, 3.0 h; yield 77%; yellow oil; $R_f = 0.38$ (CHCl₃-MeOH 95:5 v/v); IR (CHCl₃) ν 1674 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.45-7.39 (5H, m, arom. H), 5.33 (1H, s, H-6a), 2.67-2.49 (4H, m, H-2', H-6'), 2.40 (3H, s, Me), 2.34 (2H, t, $J = 7.6$ Hz, NCH₂Et), 2.29 (3H, s, Ac), 2.02-1.86 (4H, m, H-3', H-5'), 1.54-1.49 (2H, sex, $J = 7.6$ Hz, NCH₂CH₂CH₃), 0.90 (3H, t, $J = 7.6$ Hz, NC₂H₄CH₃); ¹³C NMR (125 MHz, CDCl₃) δ 194.6 (C=O), 171.1 (C-5), 137.4 (arom. C), 129.3, 128.6 (2C) (arom. CH), 125.0 (arom. CH), 114.7 (C-3a), 114.0 (C-6), 113.0 (C-2), 88.8 (C-6a), 60.1 (NCH₂CH₂), 51.2 (C-2', C-6'), 36.9, 36.7 (C-3', C-5'), 29.5 (Ac), 20.4 (NCH₂CH₂), 15.3 (Me), 11.9 (NC₂H₄CH₃); FAB HRMS (acetone/NBA): calcd for C₂₁H₂₈O₄N 358.2018 (M+H); Found 358.2027.




6-Acetyl-1'-benzyl-5-methyl-3a-phenyl-3a,6a-dihydrospiro[furo[2,3-*d*][1,3]dioxole-2,4'-piperidine]

(3af): Reaction time, 4.0 h; yield 89%; orange oil; $R_f = 0.34$ (CHCl₃-MeOH 95:5 v/v); IR (CHCl₃) ν 1674 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.45-7.37 (5H, m, arom. H), 7.31-7.28 (4H, m, arom. H), 7.25-7.22 (1H, m, arom. H), 5.32 (1H, s, H-6a), 3.50 (2H, s, NBn), 2.64-2.50 (4H, m, 2×NCH₂), 2.37 (3H, s, Me), 2.28 (3H, s, Ac), 1.99-1.83 (4H, m, 2×CH₂); ¹³C NMR (125 MHz, CDCl₃) δ 194.5 (C=O), 171.0

(C-5), 138.4, 137.4 (arom. C), 129.3, 128.6 (2C) (arom. CH), 125.0 (arom. CH), 114.7 (C-3a), 114.0 (C-6), 113.0 (C-2), 88.7 (C-6a), 62.4 (NCH₂Ph), 51.0 (2×NCH₂), 36.8, 36.6 (2×CH₂), 29.4 (Ac), 15.2 (Me); FAB HRMS (acetone/NBA): calcd for C₂₅H₂₈O₄N 406.2018 (M+H); Found 406.2032.




1',6-Diacetyl-5-methyl-3a-phenyl-3a,6a-dihydrospiro[furo[2,3-*d*][1,3]dioxole-2,4'-piperidine] (3ag): Reaction time, 2.5 h; yield quant; yellow microcrystals (from Et₂O/hexane); mp 126 °C; *R_f* = 0.28 (CHCl₃-MeOH 95:5 v/v); IR (KBr) ν 1676 (C=O), 1636 (NC=O); ¹H NMR (500 MHz, CDCl₃) δ 7.44-7.41 (5H, m, arom. H), 5.38 (1H, s, H-6a), 3.83-3.80 (1H, m, NCH_a), 3.71 (1H, t, *J* = 5.9 Hz, NCH_b), 3.65 (1H, t, *J* = 5.9 Hz, NCH_{a'}), 3.57-3.50 (1H, m, NCH_{b'}), 2.41 (3H, s, Me), 2.29 (3H, s, Ac), 2.13, 2.11 (3H, s, NAc, 1:1.3), 1.98-1.77 (4H, m, 2×CH₂); ¹H NMR (500 MHz, DMSO-*d*₆ at 80 °C) δ 7.49-7.43 (5H, m, arom. H), 5.51 (1H, s, H-6a), 3.72-3.43 (4H, m, 2×NCH₂), 2.34 (3H, s, Me), 2.23 (3H, s, Ac), 2.01 (3H, s, NAc), 1.97-1.63 (4H, m, 2×CH₂); ¹³C NMR (125 MHz, CDCl₃) δ 194.3, 194.1 (C=O), 171.1 (C-5), 168.8, 168.7 (NC=O), 136.9, 136.8 (arom. C), 129.54, 129.52 (arom. CH), 128.68, 128.66 (2C) (arom. CH), 124.9 (2C) (arom. CH), 114.9, 114.8 (C-3a), 114.1, 114.0 (C-6), 112.23, 112.20 (C-2), 88.99, 88.95 (C-6a), 44.2, 44.1 (NCH₂), 39.4, 39.3 (NC'H₂), 37.5, 37.2 (CH₂), 36.7, 36.4 (C'H₂), 29.41, 29.39 (Ac), 21.3 (NAc), 15.28, 15.25 (Me); ¹³C NMR (125 MHz, DMSO-*d*₆ at 80 °C) δ 194.5 (C=O), 171.1 (C-5), 169.4 (NC=O), 138.3 (arom. C), 130.6 (arom. CH), 129.8 (2C) (arom. CH), 126.3 (2C) (arom. CH), 115.6 (C-3a), 115.3 (C-6), 113.2 (C-2), 89.4 (C-6a), 30.4 (Ac), 22.2 (NAc), 16.0 (Me); FAB HRMS (acetone/NBA): calcd for C₂₀H₂₄O₅N 358.1654 (M+H); Found 358.1665.




6-Acetyl-1'-benzoyl-5-methyl-3a-phenyl-3a,6a-dihydrospiro[furo[2,3-*d*][1,3]dioxole-2,4'-piperidine (3ah): Reaction time, 3.0 h; yield 94%; yellow microcrystals (from Et₂O/hexane); mp 64-65 °C; *R_f* = 0.37 (CHCl₃-MeOH 95:5 v/v); IR (KBr) ν 1684 (C=C-C=O), 1653 (NC=O), 1558 (arom); ¹H NMR (500 MHz, CDCl₃) δ 7.56-7.35 (10H, m, arom. H), 5.59 (1H, s, H-6a), 4.18-3.77 (2H, m, NCH₂), 3.71-3.37 (2H, m, NC'H₂), 2.42, 2.36 (3H, s, Me, 1:1.3), 2.28 (3H, s, Ac), 2.14-2.00 (2H, m, CH₂), 1.99-1.69 (2H, m, C'H₂); ¹H NMR (500 MHz, DMSO-*d*₆ at 80 °C) δ 7.48-7.43 (10H, m, arom. H), 5.53 (1H, s, H-6a), 3.79-3.42 (4H, m, 2×NCH₂), 2.34 (3H, s, Me), 2.25 (3H, s, Ac), 1.97-1.63 (4H, m, 2×CH₂); ¹³C NMR (125 MHz, CDCl₃) δ 194.0 (C=O), 170.9 (C-5), 170.2 (NC=O), 136.7, 135.5 (arom. C), 129.6 (arom. CH), 129.4 (arom. CH), 128.5 (2C) (arom. CH), 128.3 (2C) (arom. CH), 126.6 (2C) (arom. CH), 124.8 (2C) (arom.

CH), 114.7 (C-3a), 113.9 (C-6), 112.1 (C-2), 88.9 (C-6a), 45.3 (NCH₂), 39.9 (NC'H₂), 37.5 (CH₂), 36.3 (C'H₂), 29.3 (Ac), 15.1 (Me); ¹³C NMR (125 MHz, DMSO-*d*₆ at 80 °C) δ 193.6 (C=O), 170.3 (C-5), 169.6 (NC=O), 137.4 (arom. C), 136.5 (arom. C), 129.7 (arom. CH), 129.0 (arom. CH), 128.7 (4C) (arom. CH), 127.1 (2C) (arom. CH), 125.4 (2C) (arom. CH), 114.7 (C-3a), 114.4 (C-6), 112.3 (C-2), 88.6 (C-6a), 37.1 (CH₂), 36.7 (C'H₂), 29.5 (Ac), 15.1 (Me); FAB HRMS (acetone/NBA): calcd for C₂₅H₂₆O₅N 420.1811 (M+H); Found 420.1822.




Benzyl 6-Acetyl-5-methyl-3a-phenyl-3a,6a-dihydrospiro[furo[2,3-*d*][1,3]-dioxole-2,4'-piperidine]-1'-carboxylate (3ai): Reaction time, 3.0 h; yield 94%; pale yellow oil; *R*_f = 0.36 (EtOAc-hexane 3:7 *v/v*); IR (CHCl₃) ν 1682 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.44-7.32 (10H, m, arom. H), 5.36 (1H, s, H-6a), 5.14 (2H, s, OCH₂Ph), 3.72 (2H, t, *J* = 5.8 Hz, NCH₂), 3.63-3.59 (2H, m, NC'H₂), 2.40 (3H, s, Me), 2.28 (3H, s, Ac), 1.93 (2H, m, CH₂), 1.82 (2H, m, C'H₂); ¹³C NMR (125 MHz, CDCl₃) δ 194.3 (C=O), 171.1 (C-5), 155.0 (NC=O), 136.9 (arom. C), 136.6 (arom. C), 129.5 (arom. CH), 128.6 (arom. CH), 128.4 (arom. CH), 128.0 (arom. CH), 127.8 (arom. CH), 124.9 (arom. CH), 114.8, 114.0, 112.4 (C-3a, C-2, C-6), 88.9 (C-6a), 67.2 (CH₂Ph), 41.9 (NCH₂), 41.8 (NC'H₂), 36.9 (CH₂), 36.6 (C'H₂), 29.4 (Ac), 15.2 (Me); FAB HRMS (acetone/NBA): calcd for C₂₆H₂₈O₆N 450.1917 (M+H); Found 450.1911.

Reaction of 1a with 4-Hydroxy-2*H*-chromen-2-one (5) in the Presence of BF₃•Et₂O. To a 10 mL two-necked flask, 4-hydroxy-2*H*-chromen-2-one (5) (145.6 mg, 0.9 mmol), BF₃•Et₂O (38 μL, 0.3 mmol) and dry chlorobenzene (0.5 mL) were added under argon and the mixture was heated at 130 °C. After boiling, a solution of pentenedione 1a (79.0 mg, 0.3 mmol) in dry chlorobenzene (0.3 mL) was dropwise added over 10 min. The reaction mixture was continued to heat at 130 °C until the reaction was completed. Water (30 mL) was added to the mixture and the aqueous solution was extracted with CHCl₃ (3×20 mL). The combined extracts were dried over anhydrous sodium sulfate, then concentrated to dryness. The residue was purified by column chromatography on silica gel eluting with DCM, giving furfurylchromenone 6 (101.5 mg, 77%) and furochromenone 7 (17.3 mg, 13%) (Table 3, Entry 6).




3-((3-Acetyl-5-phenylfuran-2-yl)methyl)-4-hydroxy-2H-chromen-2-one (6): Reaction time, 3.0 min; yield 77%; colorless needles (from CHCl₃/hexane); mp 241-242 °C; $R_f = 0.29$ (EtOAc-hexane 3:7 v/v); IR (KBr) ν 3482-2720 (OH), 1701 (OC=O), 1653 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 10.9 (1H, s, OH), 7.87 (1H, dd, $J = 7.9, 1.8$ Hz, H-5), 7.69 (2H, dd, $J = 7.5, 2.0$ Hz, arom, H), 7.48 (1H, td, $J = 7.9, 1.8$ Hz, H-7), 7.36 (2H, t, $J = 7.5$ Hz, arom, H), 7.30-7.22 (3H, m, H-6, H-8, arom, H), 6.79 (1H, s, H-4'), 4.29 (2H, s, CH₂), 2.62 (3H, s, Me); ¹³C NMR (125 MHz, CDCl₃) δ 200.5 (C=O), 163.6 (OC=O), 162.4 (C-4), 158.3 (C-2'), 153.6 (C-5'), 152.6 (C-8a), 132.0 (arom. CH), 128.9 (arom. C), 128.8 (arom. CH), 128.5 (arom. CH), 124.1 (arom. CH), 123.8 (arom. CH), 123.7 (arom. CH), 122.8 (C-3'), 116.3 (arom. CH), 116.2 (C-4a), 103.7 (C-4'), 99.2 (C-3), 28.9 (Me), 22.9 (CH₂); Anal. Calcd for C₂₁H₁₉O₃N: C, 73.33; H, 4.48. Found: C, 73.00; H, 4.53.




(Z)-2-(2-Hydroxy-4-oxopent-2-en-3-yl)-3-phenyl-4H-furo[3,2-c]chromen-4-one (7): Reaction time, 3.0 min; yield 13%; colorless microcrystals (from Et₂O/hexane); mp 174 °C; $R_f = 0.62$ (EtOAc-hexane 3:7 v/v); IR (KBr) ν 3500-3200 (OH), 1744 (OC=O), 1632 (C=C-C=O); ¹H NMR (500 MHz, CDCl₃) δ 16.99 (1H, s, OH), 7.92 (1H, d, $J = 6.5$ Hz, H-9), 7.57 (1H, t, $J = 7.3$ Hz, H-7), 7.49-7.36 (7H, m, H-6, H-8, arom, H), 1.96 (6H, s, 2×Me); ¹³C NMR (125 MHz, CDCl₃) δ 193.9 (C=O), 157.8 (C-10), 157.4 (OC=O), 152.7 (C-5a), 149.0 (C-2), 131.1 (C-7), 129.4 (arom. C), 129.2 (2C) (arom. CH), 128.5 (2C) (arom. CH), 128.3 (arom. CH), 124.8 (C-3), 124.5 (C-8), 120.9 (C-9), 117.3 (C-6), 112.6 (C-9a), 109.4 (C-3a), 103.6 (C-3'), 24.0 (Me); Anal. Calcd for FAB HRMS (acetone/NBA): calcd for C₂₂H₁₇O₅ 361.1076 (M+H); Found 361.1061.

Michael Addition of Pentenedione 1a with Indole. A solution of **1a** (214.9 mg, 1.0 mmol) and indole (118.3 mg, 1.0 mmol) in MeCN (1.0 mL) was heated under reflux for 9.5 h. The solvent was then removed under reduced pressure and the residue was purified by column chromatography on silica gel eluting with EtOAc/hexane (3:7 v/v), giving the desired Michael adduct **8** (136.8 mg, 97%). The reaction was also carried out in MeCN (3.0 mL) for 19 h to give **8** (98%); DCM for 24 h, **8** (93%); MeOH for 14 h, **8** (80%) along with **12** (16%); THF for 48 h, **8** (40%); DMF for 3 h, **8** (83%) and **12** (8%); DCE for 36 h, **8** (98%); and toluene for 23 h, **8** (97%) (Scheme 8).



3-Acetyl-2-(1*H*-indol-3-yl)-1-phenylpentane-1,4-dione (8): Reaction time, 9.5 h; yield 97%; colorless microcrystals (from benzene); mp 157 °C; R_f = 0.20 (EtOAc-hexane 2:8 v/v); IR (KBr) ν 3500–3200 (NH), 1717, 1705, 1676 (C=O); ^1H NMR (500 MHz, CDCl_3) δ 8.34 (1H, br s, NH), 7.95 (2H, dd, J = 8.0, 1.3 Hz, arom. H), 7.74–7.72 (1H, m, H-4), 7.39 (1H, t, J = 8.0 Hz, arom. H), 7.30–7.25 (3H, m, H-7, arom. H), 7.18–7.13 (2H, m, H-5, H-6), 7.03 (1H, d, J = 3.0 Hz, H-2), 5.68 (1H, d, J = 10.5 Hz, H-2'), 5.05 (1H, d, J = 10.5 Hz, H-3'), 2.30 (3H, s, Ac), 1.88 (3H, s, Ac); ^{13}C NMR (125 MHz, CDCl_3) δ 203.7, 202.0 (MeC=O), 197.7 (PhC=O), 136.3 (arom C), 135.7 (C-7a), 133.0 (arom. CH), 128.7 (2C) (arom. CH), 128.4 (2C) (arom. CH), 125.7 (C-3a), 123.9 (C-2), 122.6 (C-5), 120.4 (C-6), 119.0 (C-4), 111.5 (C-7), 109.3 (C-3), 70.2 (C-3'), 45.3 (C-2'), 31.4, 30.2 (2×Me); Anal. Calcd for $\text{C}_{21}\text{H}_{19}\text{O}_3\text{N}$: C, 75.66; H, 5.74; N, 4.20. Found: C, 75.72; H, 5.79; N, 4.10.


Michael Addition of Pentenedione 1a with Pyrrole. A solution of **1a** (108.3 mg, 0.5 mmol) and pyrrole (102.0 mg, 1.5 mmol) in MeCN (0.5 mL) was heated under reflux for 3.5 h. The solvent was then removed in vacuo and the residue was purified by column chromatography on silica gel eluting with EtOAc/hexane (2:8-4:6 v/v), giving the desired monoalkylpyrrole **9** (122.7 mg, 87%) (Scheme 8).



3-Acetyl-1-phenyl-2-(1*H*-pyrrol-2-yl)pentane-1,4-dione (9): Reaction time, 3.5 h; yield 84%; colorless microcrystals (from Et_2O); mp 115–116 °C; R_f = 0.43 (EtOAc-hexane 3:7 v/v); IR (KBr) ν 3500–3160 (NH), 1724, 1701, 1663 (C=O); ^1H NMR (500 MHz, CDCl_3) δ 8.38 (1H, br s, NH), 7.96 (2H, dd, J = 7.4, 1.2 Hz, arom. H), 7.51 (1H, tt, J = 7.4, 1.2 Hz, arom. H), 7.40 (2H, t, J = 7.4 Hz, arom. H), 6.69 (1H, dt, J = 5.0, 2.1 Hz, H-5), 6.06 (1H, dd, J = 5.0, 2.1 Hz, H-3), 6.00 (1H, td, J = 5.0, 1.5 Hz, H-4), 5.47 (1H, d, J = 11 Hz, H-2'), 4.86 (1H, d, J = 11 Hz, H-3'), 2.24 (3H, s, Ac), 2.02 (3H, s, Ac); ^{13}C NMR (125 MHz, CDCl_3) δ 204.1, 201.4 (MeC=O), 197.3 (PhC=O), 135.7 (arom. C), 133.4 (arom. CH), 128.8 (arom. CH),


128.6 (arom. CH), 123.7 (C-2), 119.1 (C-5), 109.3 (C-4), 108.8 (C-3), 70.1 (C-3'), 46.6 (C-2'), 31.6, 30.1 (2×Me); FAB HRMS (acetone/NBA): calcd for C₁₇H₁₇O₃N 283.1208 (M⁺); Found 283.1205.

To a 10-mL round-bottomed flask, **1a** (230.6 mg, 1.0 mmol), pyrrole (35.7 mg, 0.5 mmol), and MeCN (0.5 mL) were added, and the mixture was heated under reflux for 1 h. After work-up as previously described, bis(alkyl)pyrrole was obtained (208.7 mg; 79% yield).




2,2'-(1H-Pyrrole-2,5-diyl)bis(3-acetyl-1-phenylpentane-1,4-dione): Pink microcrystals (from Et₂O); mp 240 °C; *R*_f = 0.20 (EtOAc-hexane 3:7 v/v); IR (KBr) ν 3400–3200 (NH), 1734, 1684, 1653 (C=O), 1558 (arom); ¹H NMR (500 MHz, CDCl₃) δ 8.56 (1H, br s, NH), 7.89 (4H, dd, *J* = 8.5, 1.0 Hz, arom, H), 7.46–7.43 (2H, m, arom. H), 7.30 (4H, t, *J* = 8.0 Hz, arom. H), 5.80 (2H, d, *J* = 2.5 Hz, H-3, H-4), 5.41 (2H, d, *J* = 10.8 Hz, 2×H-2'), 4.75 (2H, d, *J* = 10.8 Hz, 2×H-3'), 2.23 (6H, s, 2×Ac), 1.83 (6H, s, 2×Ac); ¹³C NMR (125 MHz, CDCl₃) δ 204.0, 201.2 (MeC=O), 196.4 (PhC=O), 135.5 (arom. C), 133.1 (arom. CH), 128.8 (arom. CH), 128.4 (arom. CH), 125.8 (C-2, C-5), 109.6 (C-3, C-4), 69.5 (C-3'), 47.1 (C-2'), 32.0, 30.3 (2×Me); FAB HRMS (acetone/NBA): calcd for C₃₀H₂₉O₆N 499.1995 (M⁺); Found 499.2004.

Michael Addition of Pentenedione 1a with Furan. A solution of **1a** (108.0 mg, 0.5 mmol) and furan (362 μ L, 5.0 mmol) in AcOH (0.5 mL) was stirred at room temperature for 4.0 h. The solvent was then removed under reduced pressure and the residue was purified by column chromatography on silica gel eluting with EtOAc/hexane (1:3 v/v), giving the desired alkylfuran **10** (107.4 mg, 76%) (Scheme 8).




3-Acetyl-2-(furan-2-yl)-1-phenylpentane-1,4-dione (10): Reaction time, 4.0 h; yield 76%; colorless prisms (from Et₂O); mp 95–96 °C; *R*_f = 0.47 (EtOAc-hexane 3:7 v/v); IR (KBr) ν 1734, 1701, 1653 (C=O), 1558 (arom); ¹H NMR (500 MHz, CDCl₃) δ 7.98 (2H, d, *J* = 7.5 Hz, arom. H), 7.51 (1H, t, *J* = 7.5 Hz, arom. H), 7.40 (2H, t, *J* = 7.8 Hz, arom. H), 7.30 (1H, d, *J* = 1.7 Hz, H-5), 6.25 (1H, dd, *J* = 3.2, 1.7 Hz, H-4), 6.20 (1H, d, *J* = 3.2 Hz, H-3), 5.56 (1H, d, *J* = 11.3 Hz, H-2'), 4.92 (1H, d, *J* = 11.3 Hz, H-3'), 2.29 (3H, s, Ac), 2.09 (3H, s, Ac); ¹³C NMR (125 MHz, CDCl₃) δ 202.5, 201.0 (MeC=O), 194.7 (PhC=O), 148.3 (C-2), 142.9 (C-5), 135.3 (arom. C), 133.4 (arom. CH), 128.8 (arom. CH), 128.5 (arom. CH), 111.1 (C-4), 109.3 (C-3), 68.6 (C-3'), 47.0 (C-2'), 30.3, 30.0 (2×Me); FAB HRMS (acetone/NBA): calcd for C₁₇H₁₇O₄ 285.1127 (M⁺H); Found 285.1137.

Michael Addition of Pentenedione **1a with *N*-Methylaniline.** A solution of **1a** (107.5 mg, 0.5 mmol) and *N*-methylaniline (165 μ L, 1.5 mmol) in AcOH (0.5 mL) was heated under reflux for 6.0 min. The solvent was then removed under reduced pressure and the residue was purified by thin layer chromatography on silica gel eluting with CHCl₃/MeOH (98:2 v/v), giving the desired adduct **11** (120.3 mg, 75%) (Scheme 8).



3-Acetyl-2-(4-(methylamino)phenyl)-1-phenylpentane-1,4-dione (11**):** Reaction time, 6.0 min; yield 75%; pale yellow amorphous; R_f = 0.38 (CHCl₃-MeOH 98:2 v/v); IR (CH₃Cl) ν 3474-3360 (NH), 1730, 1697, 1676 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.93 (2H, dd, J = 7.8, 1.0 Hz, arom. H), 7.43 (1H, tt, J = 7.8, 1.0 Hz, arom. H), 7.34 (2H, t, J = 7.8 Hz, arom. H), 7.05 (2H, dd, J = 6.6, 1.9 Hz, H-3), 6.49 (2H, dd, J = 6.6, 1.9 Hz, H-2), 5.24 (1H, d, J = 11.5 Hz, H-2'), 4.81 (1H, d, J = 11.5 Hz, H-3'), 3.72 (1H, br s, NH), 2.75 (3H, s, NMe), 2.26 (3H, s, Ac), 1.95 (3H, s, Ac); ¹³C NMR (125 MHz, CDCl₃) δ 203.7, 201.7 (2 \times MeC=O), 197.9 (PhC=O), 148.8 (C-1), 135.8 (arom. C), 132.8 (arom. CH), 129.5 (arom. CH), 128.8 (arom. CH), 128.3 (arom. C), 122.4 (C-4), 112.9 (C-2), 71.1 (C-3'), 53.0 (C-2'), 31.7 (NMe), 30.3, 30.1 (2 \times Me); FAB HRMS (acetone/NBA): calcd for C₂₀H₂₁O₃N 323.1521 (M⁺); Found 323.1529.


Synthesis of Furyl-substituted Indole **12.** To a 10-mL two-necked flask, pentenedione **1a** (108.3 mg, 0.5 mmol), indole (70.1 mg, 0.6 mmol), BF₃·Et₂O (32 μ g, 0.25 mmol) and DCM (1.5 mL) were added and the mixture was heated under reflux for 30 min. After the reaction was completed, water (10 mL), brine (10 mL), and a saturated aqueous solution of NaHCO₃ (30 mL) were then added, and the aqueous mixture was extracted with CHCl₃ (3 \times 20 mL). The combined extracts were washed with water (20 mL), dried over anhydrous sodium sulfate, then concentrated to dryness. The residue was separated by column chromatography on silica gel eluting with EtOAc/hexane (2:8 v/v), giving the desired furyl-substituted indole **12** (138.9 mg, 88%) (Scheme 9).



3-(3-Acetyl-2-methyl-5-phenylfuran-4-yl)-1H-indole (12**):**^{10c} Yellow cubes (from DCM); mp 141 $^{\circ}$ C; R_f = 0.47 (EtOAc-hexane 3:7 v/v); IR (KBr) ν 3400–3200 (NH), 1653 (C=O), 1558 (arom); ¹H NMR (500 MHz, CDCl₃) δ 8.63 (1H, br s, NH), 7.43-7.38 (4H, m, arom, H), 7.23 (1H, t, J = 7.0 Hz, arom. H),

7.14-7.08 (5H, m, arom. H), 2.69 (3H, s, Me), 1.85 (3H, s, Ac); ^{13}C NMR (125 MHz, CDCl_3) δ 196.9 (C=O), 157.6 (C-5'), 148.5 (C-2'), 136.1 (C-7a), 130.5 (arom, C), 128.2 (arom, CH), 127.7 (C-3a), 127.2 (arom, CH), 125.3 (C-4'), 125.1 (arom. CH), 123.7 (arom. CH), 122.6 (arom. CH), 120.4 (arom. CH), 119.7 (arom, CH), 113.2 (C-3'), 111.4 (arom, CH), 108.3 (C-3), 29.9 (Ac), 14.7 (Me).

Paal-Knorr Reaction of Alkylindole **8 to Pyrrolyindole **13**.** To a 10-mL round-bottomed flask, alkylindole **8** (128.9 mg, 0.4 mmol), ammonium acetate (66.2 mg, 0.8 mmol) and AcOH (0.3 mL) were added and the mixture was heated at 110 °C for 45 min. After the reaction was completed, water (10 mL) was added to the reaction mixture. The resulting aqueous mixture was then extracted with CHCl_3 (3×20 mL) and the combined extracts were washed with a saturated aqueous solution of NaHCO_3 (20 mL), water (20 mL), dried over anhydrous sodium sulfate, then concentrated to dryness, giving the desired pyrrole-substituted indole **13a** (125.0 mg, quant) without purification (Scheme 9).




3-(3-Acetyl-2-methyl-5-phenyl-1H-pyrrol-4-yl)-1H-indole (13a): Reaction time, 45 min; yield quant; yellow microcrystals (from DCM); mp 96 °C; R_f = 0.33 (EtOAc-hexane 3:7 v/v); IR (KBr) ν 3500-2500 (NH), 1636 (C=O); ^1H NMR (500 MHz, CDCl_3) δ 9.13 (1H, br s, pyrrole-NH), 8.45 (1H, br s, indole-NH), 7.41 (1H, d, J = 7.8 Hz, H-4), 7.36 (1H, d, J = 7.8 Hz, H-7), 7.18 (1H, t, J = 7.8 Hz, H-6), 7.15-7.13 (2H, m, arom. H), 7.07-7.03 (4H, m, arom. H), 6.94 (1H, d, J = 2.5 Hz, H-2), 2.60 (3H, s, Me), 1.84 (3H, s, Ac); ^{13}C NMR (125 MHz, CDCl_3) δ 198.0 (C=O), 136.1 (C-7a), 135.9 (C-2'), 132.4 (arom. C), 128.9 (C-3a), 128.6 (C-5'), 128.3 (2C) (arom. CH), 126.3 (3C) (arom. CH), 123.7 (C-2), 123.6 (C-3'), 122.1 (C-5), 119.97, 119.94 (C-4, C-6), 114.0 (C-4'), 111.2 (C-7), 111.1 (C-3), 29.7 (O=CMe), 14.6 (Me); FAB HRMS (acetone/NBA): calcd for $\text{C}_{21}\text{H}_{18}\text{ON}_2$ 314.1419 (M⁺); Found 314.1427.

The alkylindole **8** (99.8 mg, 0.3 mmol) and benzylamine (66 mL, 0.6 mmol) were heated in AcOH (0.3 mL) at 110 °C for 20 min. After the previously described work-up, the residue was purified by column chromatography on silica gel eluting with EtOAc/hexane (4:6 v/v), giving the desired pyrrole **13b** (107.4 mg, 89%) (Scheme 9).

NH), 7.27 (1H, d, $J = 7.8$ Hz, H-4), 7.24-7.20 (5H, m, arom, H), 7.16 (1H, d, $J = 7.8$ Hz, H-7), 7.04 (1H, d, $J = 2.5$ Hz, H-2), 7.50 (1H, td, $J = 7.8, 1.0$ Hz, H-6), 6.89 (1H, td, $J = 7.8, 1.5$ Hz, H-5), 4.90 (1H, t, $J = 5.5$ Hz, OH), 3.87 (2H, t, $J = 7.4$ Hz, CH_2OH), 3.39 (2H, td, $J = 7.4, 4.6$ Hz, NCH_2), 2.57 (3H, s, Me), 1.66 (3H, s, Ac); ^{13}C NMR (125 MHz, $\text{DMSO-}d_6$) δ 195.6 (C=O), 135.6 (C-7a), 134.5 (C-2'), 132.2 (arom. C), 131.9 (C-5'), 131.2 (2C) (arom. CH), 128.6 (C-3a), 127.8 (2C) (arom. CH), 127.5 (arom. CH), 124.7 (C-2), 121.8 (C-3'), 120.8 (C-5), 118.8, 118.7 (C-4, C-6), 114.8 (C-4'), 111.4 (C-7), 109.6 (C-3), 60.0 (CH_2OH), 45.8 (NCH_2), 29.5 (O=CMe), 12.0 (Me); FAB HRMS (acetone/NBA): calcd for $\text{C}_{23}\text{H}_{22}\text{O}_2\text{N}_2$ 358.1681 (M⁺); Found 358.1689.

Paal-Knorr Reaction of 8 to Thienylindole 14. A mixture of **8** (66.3 mg, 0.2 mmol) and Lawesson's reagent (123.1 mg, 0.3 mmol) was heated in toluene (3.0 mL) at 50 °C for 30 min under argon. After the reaction was completed, water (10 mL) was added. The resulting aqueous mixture was extracted with CHCl_3 (3×20 mL) and the combined extracts were dried over anhydrous sodium sulfate, then concentrated to dryness. The obtained residue was separated by column chromatography on silica gel eluting with EtOAc/hexane (2:8 v/v), giving the desired thienylindole **14** (32.5 mg, 45%) together with furylindole **12** (14.2 mg, 23%) (Scheme 9).



3-(3-Acetyl-2-methyl-5-phenylthiophen-4-yl)-1H-indole (14): Reaction time, 30 min; yield 45%; yellow microcrystals (from DCM); mp 159 °C; $R_f = 0.12$ (EtOAc-hexane 3:7 v/v); IR (KBr) ν 3500-3120 (NH), 1655 (C=O); ^1H NMR (500 MHz, CDCl_3) δ 8.29 (1H, br s, NH), 7.35 (1H, d, $J = 8.0$ Hz, H-4), 7.30 (1H, d, $J = 8.0$ Hz, H-7), 7.19-7.11 (6H, m, H-6, arom. H), 7.02 (1H, t, $J = 7.0$ Hz, H-5), 6.95 (1H, d, $J = 2.5$ Hz, H-2), 2.62 (3H, s, Me), 1.82 (3H, s, Ac); ^{13}C NMR (125 MHz, CDCl_3) δ 200.2 (C=O), 142.5 (C-2'), 141.3 (C-3'), 136.9 (C-4'), 135.8 (C-7a), 134.1 (arom. C), 129.4 (C-5'), 128.6 (2C) (arom. CH), 128.2 (2C) (arom. CH), 127.4 (C-3a), 127.1 (arom. CH), 123.7 (C-2), 122.5 (C-6), 120.3 (C-5), 119.9 (C-4), 111.8 (C-3), 111.1 (C-7), 30.4 (O=CMe), 15.1 (Me); FAB HRMS (acetone/NBA): calcd for $\text{C}_{21}\text{H}_{17}\text{ONS}$ 331.1031 (M⁺); Found 331.1031.

ACKNOWLEDGEMENTS

This research was supported by a Grant-in-Aid for Scientific Research (C), No. 25410049, from the Japan Society for the Promotion of Science. We also acknowledge Nissan Chemical Industries, Ltd., and Astellas Foundation for Research on Metabolic Disorders for their financial support.

REFERENCES AND NOTES

1. a) J. A. Joule and K. Mills, 'Heterocyclic Chemistry at a Glance,' Wiley, Chichester, 2012; b) J. J. Li and E. J. Corey, 'Name Reactions in Heterocyclic Chemistry,' Wiley-VCH, Weinheim, 2005; c) J. A. Joule and K. Mills, 'Heterocyclic Chemistry,' 5th ed, Wiley-Blackwell, Chichester, 2010.
2. a) S. Ren, S. Song, L. Ye, C. Feng, and T.-P. Loh, *Chem. Commun.*, 2016, **52**, 10373; b) S. Xing, J. Ren, K. Wang, H. Cui, T. Xia, M. Zhang, and D. Wang, *Adv. Synth. Catal.*, 2016, **358**, 3093; c) N. V. Chechina, V. V. Zubar, I. V. Omelchenko, and N. N. Kolos, *ARKIVOC*, 2015, **vii**, 293; d) Y. Yang, F. Ni, W.-M. Shu, and A.-X. Wu, *Tetrahedron*, 2014, **70**, 6733; e) W.-M. Shu, Y. Yang, D.-X. Zhang, L.-M. Wu, Y.-P. Zhu, G.-D. Yin, and A.-X. Wu, *Org. Lett.*, 2013, **15**, 456; f) G. Blay, I. Fernández, A. Monleón, J. R. Pedro, and C. Vila, *Tetrahedron*, 2009, **65**, 9264; g) H. Quast, M. Seefelder, S. Ivanova, M. Heubes, E.-M. Peters, and K. Peters, *Eur. J. Org. Chem.*, 1999, 3343.
3. a) T. D. Avery, D. K. Taylor, and E. R. T. Tiekink, *J. Org. Chem.*, 2000, **65**, 5531; b) W. Adam, M. Balci, and H. Kilic, *J. Org. Chem.*, 2000, **65**, 5926.
4. a) W. Adam and A. Rodriguez, *J. Am. Chem. Soc.*, 1980, **102**, 404; b) M. L. Graziano, M. R. Iesce, and R. Scarpati, *J. Chem. Soc., Perkin Trans. 1*, 1980, 1955; c) M. L. Graziano, M. R. Iesce, B. Carli, and R. Scarpati, *Synthesis*, 1982, 736; d) M. L. Graziano, M. R. Iesce, B. Carli, and R. Scarpati, *Synthesis*, 1983, 125; e) M. L. Graziano, M. R. Iesce, S. Chiosi, and R. Scarpati, *J. Chem. Soc., Perkin Trans. 1*, 1983, 2071; f) M. L. Graziano, M. R. Iesce, and R. Scarpati, *Synthesis*, 1984, 64; g) M. L. Graziano, M. R. Iesce, and R. Scarpati, *Synthesis*, 1984, 66; h) M. R. Iesce, F. Cermola, M. L. Graziano, and R. Scarpati, *J. Chem. Soc., Perkin Trans. 1*, 1994, 147; i) M. R. Iesce, F. Cermola, A. Guitto, and F. Giordano, *J. Chem. Soc., Perkin Trans. 1*, 1999, 475.
5. a) S. Onitsuka, H. Nishino, and K. Kurosawa, *Heterocycl. Commun.*, 2000, **6**, 529; b) S. Onitsuka, H. Nishino, and K. Kurosawa, *Tetrahedron*, 2001, **57**, 6003.
6. a) F. Ni, Y. Yang, W.-M. Shu, J.-R. Ma, and A.-X. Wu, *Org. Biomol. Chem.*, 2014, **12**, 9466; b) Y. Yang, F. Ni, W.-M. Shu, and A.-X. Wu, *Chem. Eur. J.*, 2014, **20**, 11776; c) C. Asta, D. Schmidt, J. Conrad, W. Frey, and U. Beifuss, *Org. Biomol. Chem.*, 2013, **11**, 5692; d) W. Friedrichsen, 'Comprehensive Heterocyclic Chemistry II,' ed. by A. R. Katritzky, C. W. Rees, and E. F. V. Scriven, Pergamon, Oxford, 1996, Vol. 2, pp. 351-393.
7. a) S. Onitsuka, H. Nishino, and K. Kurosawa, *Tetrahedron Lett.*, 2000, **41**, 3149; b) S. Onitsuka and H. Nishino, *Tetrahedron*, 2003, **59**, 755.

8. X-Ray coordinates were deposited with the Cambridge Crystallographic Data Centre (<https://www.ccdc.cam.ac.uk>), deposition number: CCDC 1546190.
9. a) S. Dey, D. Nandi, P. K. Pradhan, V. S. Giri, and P. Jaisankar, *Tetrahedron Lett.*, 2007, **48**, 2573; b) M. Shibasaki, H. Sasai, and T. Arai, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1236; c) M. Yamaguchi, T. Shiraishi, and M. Hirama, *J. Org. Chem.*, 1996, **61**, 3520; d) Y. Zhou, Z.-L. Xia, Q. Gu, and S.-L. You, *Org. Lett.*, 2017, **19**, 762; e) S.-W. Li, J. Gong, and Q. Kang, *Org. Lett.*, 2017, **19**, 1350; f) A. V. Aksenov, N. A. Aksenov, A. A. Skomorokhov, I. V. Aksenova, G. D. Gryaznov, L. G. Voskressensky, and M. A. Rubin, *Chem. Heterocycl. Compd.*, 2016, **52**, 923; g) Y. Yu, Y. Chen, W. Wu, and H. Jiang, *Chem. Commun.*, 2017, **53**, 640.
10. a) G. Yin, Z. Wang, A. Chen, M. Gao, A. Wu, and Y. Pan, *J. Org. Chem.*, 2008, **73**, 3377; b) A. Rahmatpour and J. Aalaie, *Heteroat. Chem.*, 2011, **22**, 85; c) Y. Yang, M. Gao, L.-M. Wu, C. Deng, D.-X. Zhang, Y. Gao, Y.-P. Zhu, and A.-X. Wu, *Tetrahedron*, 2011, **67**, 5142; d) Y. Ju, D. Miao, R. Yu, and S. Koo, *Org. Biomol. Chem.*, 2015, **13**, 2588.
11. It took 8 h until the reaction in the presence of methanesulfonic acid was finished in boiling MeCN, giving **12** (96% yield).^{10c}
12. a) P. Diana, A. Carbone, P. Barraja, A. Montalbano, A. Martorana, G. Dattolo, O. Gia, L. Dalla Via, and G. Cirrincione, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 2342; b) M. M. M. Raposo, A. M. B. A. Sampaio, and G. Kirsch, *Synthesis*, 2005, 199.
13. a) I. Cobo, M. I. Matheu, S. Castellón, O. Boutureira, and B. G. Davis, *Org. Lett.*, 2012, **14**, 1728; b) F. J. L. Herrera, M. V. Fernandez, and R. G. Segura, *Carbohydr. Res.*, 1984, **127**, 217; c) J. S. Yadav, B. V. S. Reddy, D. Narsimhaswamy, P. Naga Lakshmi, K. Narsimulu, G. Srinivasulu, and A. C. Kunwar, *Tetrahedron Lett.*, 2004, **45**, 3493; d) H. Paulsen and D. Stoye, *Chem. Ber.*, 1966, **99**, 908; e) P. Hadzic and M. Popsavin, *Carbohydr. Res.*, 2010, **345**, 543; f) B. G. Roy, A. Roy, B. Achari, and S. B. Mandal, *Tetrahedron Lett.*, 2006, **47**, 7783; g) D. H. R. Barton, P. Dalko, and S. D. Géro, *Tetrahedron Lett.*, 1992, **33**, 1883; h) R. L. Dorta, C. G. Francisco, and E. Suárez, *Tetrahedron Lett.*, 1994, **35**, 2049.
14. C. O. Janssen, S. Lim, E. P. Lo, K. F. Wanb, V. C. Yu, M. A. Lee, S. B. Ng, M. J. Everett, A. D. Buss, D. P. Lane, and R. S. Boyce, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 5771.