

# Accumulation of Pericardial Fat Correlates with Left Ventricular Diastolic Dysfunction in Patients with Normal Ejection Fraction

journal or publication title	Journal of Cardiology
volume	59
number	3
page range	344-351
year	2012-05
URL	<a href="http://hdl.handle.net/2298/29229">http://hdl.handle.net/2298/29229</a>

doi: 10.1016/j.jjcc.2012.01.006

**Accumulation of Pericardial Fat Correlates with Left Ventricular Diastolic  
Dysfunction in Patients with Normal Ejection Fraction**

**First author:** Masaaki Konishi

5 **Brief title:** Pericardial fat and diastolic dysfunction

**Authors:** Masaaki Konishi, MD<sup>1,3</sup>, Seigo Sugiyama, MD<sup>1</sup>, Koichi Sugamura, MD<sup>1</sup>,  
Toshimitsu Nozaki, MD<sup>1</sup>, Junichi Matsubara, MD<sup>1</sup>, Eiichi Akiyama, MD<sup>1</sup>, Daisuke  
Utsunomiya, MD<sup>2</sup>, Yasushi Matsuzawa, MD<sup>3</sup>, Yasuyuki Yamashita, MD<sup>2</sup>, Kazuo  
Kimura, MD<sup>3</sup>, Satoshi Umemura, MD<sup>4</sup>, Hisao Ogawa, MD<sup>1</sup>

10

Departments of <sup>1</sup>Cardiovascular Medicine, <sup>2</sup>Diagnostic Radiology, Faculty of Life  
Sciences, Graduate School of Medical Sciences, Kumamoto University, Kumamoto,  
JAPAN, <sup>3</sup>Division of Cardiology, Yokohama City University Medical Center,  
Yokohama, JAPAN, <sup>4</sup>Department of Medical Science and Cardiorenal Medicine,  
15 Yokohama City University Graduate School of Medicine, Yokohama, JAPAN

**Address for correspondence:**

Seigo Sugiyama, MD, PhD

Department of Cardiovascular Medicine

20 Faculty of Life Sciences, Kumamoto University

1-1-1 Honjo, Kumamoto City 860-8556, Japan

Phone: +81-96-373-5175, Fax: +81-96-362-3256

E-mail: [ssugiyam@kumamoto-u.ac.jp](mailto:ssugiyam@kumamoto-u.ac.jp)

25 **Total word count: 4186(abstract 223)**

**Clinical trial registration:** <https://center.umin.ac.jp> ID: UMIN000001581

## Abstract

30

**Background** Left ventricular diastolic dysfunction (LVDD) plays an important role in heart failure with normal left ventricular ejection fraction (LVEF). Obesity is one of the major comorbid conditions of LVDD. Pericardial fat (PF) is an ectopic fat depot with possible paracrine or mechanical effects on the coronary circulation and myocardial function.

35

**Methods** We measured PF volume on 64 slice computed tomography and analyzed echocardiographic parameters to confirm LVDD in 229 consecutive patients suspected of coronary artery disease with LVEF of more than 50% and no symptomatic heart failure (59% men, 67±12 years). LVDD was defined as the ratio of transmitral Doppler early filling velocity to tissue Doppler early diastolic mitral annular velocity ( $E/e'$ ) >10.

40

**Results** PF volume correlated significantly with  $E/e'$  ( $r=0.21$ ,  $p<0.01$ ), left ventricular mass index ( $r=0.23$ ,  $p<0.001$ ), and left atrial diameter ( $r=0.32$ ,  $p<0.001$ ). The mean PF volume was significantly greater in patients with LVDD ( $184\pm 61$  cm<sup>3</sup>,  $n=141$ ) than in those without LVDD ( $154\pm 58$ ,  $n=88$ ,  $p<0.001$ ). Multivariate logistic regression analysis indicated that PF volume correlated significantly with the presence of LVDD (odds ratio: 2.00 per 100 cm<sup>3</sup> increase in PF volume,  $p=0.02$ ) independent of age, gender, abdominal obesity, hypertension, and diabetes.

45

**Conclusions** PF volumes are significantly associated with LVDD, independent of other factors such as hypertension or diabetes. PF may be implicated in the

50

pathogenesis of LVDD in patients with normal LVEF.

**Key words:** pericardium, obesity, diastole, heart failure

## Introduction

55

Increasing evidence suggests that approximately half of all patients with heart failure have a normal left ventricular ejection fraction (LVEF)<sup>1</sup>, and the mortality rate for this group is similar to those with heart failure and reduced LVEF<sup>2</sup>. Left ventricular diastolic dysfunction (LVDD) plays an important role in heart failure with normal ejection fraction (EF)<sup>3</sup>. However, there is no specific therapeutic strategy for LVDD partly because of the lack of understanding of its pathophysiological mechanism(s).

60

65

LVDD is most prevalent among elderly obese women with hypertension, diabetes, coronary artery disease, and/or atrial fibrillation<sup>4,5</sup>. Obesity is one of the important features of LVDD and fat tissue is known to secrete many adipokines that have local and systemic effects on the cardiovascular system<sup>6</sup>. Ectopic fat depots in muscle, liver, and pancreas are described as lipotoxic and pericardial fat (PF) is recognized as an ectopic visceral fat depot in close proximity to the myocardium and coronary arteries. PF volume is increased in obese patients and correlates with the presence<sup>7,8</sup> and incidence<sup>9</sup> of coronary atherosclerosis independent of other risk factors, indicating that PF may have paracrine or mechanical effects on the coronary circulation and myocardium. PF volume has also been reported to correlate with left atrial diameter, which is an indirect structural parameter of LVDD<sup>10-12</sup>. However, there is currently no report of a functional parameter of LVDD by tissue Doppler

70

75

echocardiography that correlates with PF volume. The purpose of this study was to assess the association between PF volume and LVDD, as determined by tissue Doppler echocardiography.

## Methods

### Study Sample

85

We analyzed consecutive inpatients in a stable condition who underwent 64-slice computed tomography (CT) coronary angiography and echocardiography between 2006 and 2009 on suspicion of coronary artery disease. None of the patients had  
90 history of previous thoracic surgery, percutaneous coronary intervention, or symptomatic heart failure. We screened 304 patients, each with LVEF more than 50%, and excluded those with previous pacemaker implantation (n=19), chronic atrial fibrillation (n=12), end-stage renal disease (n=14), significant valvular disease (n=9), poor image quality (n=17), and incomplete data (n=6). Thus, we measured PF  
95 volumes from CT images and examined echocardiographic parameters in 229 consecutive patients. The study was approved by the ethics review committee of our institution and a signed informed consent was obtained from each patient before participation. This study was registered at UMIN protocol registration system with the identification number UMIN000003361.

100



## Cardiac CT Scan Protocol

105

The 64-detector CT (Brilliance-64, Phillips Medical Systems, Cleveland, OH) was used with the following parameters: detector collimation 64 x 0.625 mm, table feed 19.7 mm/sec, 0.2 beam pitch, rotation time 420 msec, tube current 429 mA, and voltage 120 kVp, as reported previously<sup>7</sup>. Reconstructions sets at 75% of the cardiac cycle or at a particular optimal phase were prepared from the raw data files. The contrast material (Omnipaque-350; Daiichi-Sankyo Pharmaceutical, Tokyo, Japan) was administered using a mechanical power injector through a 20-gauge cannula inserted into the antecubital vein. To minimize differences in arterial enhancement across patients, we used a body-weight-tailored contrast material dose (0.7 mL/kg) and a fixed injection duration (9 sec)<sup>13</sup>. An oral  $\beta$ -blocker (metoprolol, 20 mg) was administered 1 hour prior to CT imaging, and nitroglycerin (0.3 mg) was administered immediately prior to CT imaging. The reconstructed CT image data were transferred to a workstation for post-processing (ZIO M900, Amin/ZIO, Tokyo).

120

## Cardiac CT Image Analysis

125 PF volume was measured three-dimensionally in all patients using contrast-enhanced images, as reported previously<sup>7</sup>. A predefined image display setting was used

(window width = 150 Hounsfield units (HU), window center = -120 HU) to identify pixels that correspond to fat tissue<sup>14</sup>. The readers, who were blinded to the clinical results, trimmed along the pericardial sac using axial, coronal, and sagittal slices and volume-rendered images. PF was defined to be any adipose tissue located within the pericardial sac (Figure 1). A slice 1 cm above the most cranial slice including the left anterior descending coronary artery was defined to be the superior border of the PF.

Three major coronary arteries were analyzed visually and quantitatively by contrast-enhanced coronary CT angiography. Coronary artery disease was defined to be  $\geq 75\%$  stenosis (according to the American Heart Association classification) on conventional coronary angiography (n=136) analyzed quantitatively by coronary angiography software (CAAS, Pie Medical Imaging, Maastricht, Netherlands) or  $\geq 50\%$  luminal narrowing on CT images in patients without conventional coronary angiography (n=98).

## **Echocardiography**

For each patient, echocardiography was performed by a specialized echocardiologist. The cavity dimension and wall thickness were measured in a parasternal long axis view. Left ventricular mass was estimated using the formula recommended by recent guidelines<sup>15</sup>. Measurement of LVEF was performed in biplane apical (2-and 4-chamber) views using a modified Simpson's method. The pulsed Doppler sample volume was positioned at the opened leaflet tips of mitral valve. Early and late

150 diastolic peak flow velocity and E-wave deceleration time were measured by  
transmitral Doppler imaging. The ratio of transmitral Doppler early filling velocity to  
tissue Doppler early diastolic mitral annular velocity ( $E/e'$ ) was also measured. LVDD  
was defined as the tissue Doppler oriented criterion using  $E/e' > 10^{16-18}$ .

155

### **Assessment of Risk Factors and Covariates**

Obesity was defined as a body mass index of  $\geq 25 \text{ kg/m}^2$ . Abdominal obesity was  
160 defined as a waist circumference of  $\geq 85 \text{ cm}$  for Japanese males and  $\geq 90 \text{ cm}$  for  
Japanese females<sup>19</sup>. Blood was drawn after an overnight fast. Diabetes mellitus was  
diagnosed based on the criteria set by the World Health Organization or the use of  
hypoglycemic agents or insulin. Hypertension was defined as a systolic blood pressure  
 $\geq 140 \text{ mmHg}$ , a diastolic blood pressure  $\geq 90 \text{ mmHg}$ , or the use of an antihypertensive  
165 treatment. The estimated glomerular filtration rate was calculated using a modified  
formula from the Modification of Diet in Renal Disease study equation, which was  
proposed by the Japanese Society of Nephrology<sup>20</sup>. Metabolic syndrome was  
diagnosed according to the modified Adult Treatment Panel III criteria, which include  
the waist circumference cut-off values mentioned above and the presence of three or  
170 more metabolic abnormalities.

## Statistical Analysis

175

Data are expressed as mean  $\pm$  standard deviation. Differences in risk factors and adiposity traits in patients with or without LVDD were assessed. Differences between continuous variables were analyzed by the unpaired Student's *t*-test or the Mann-Whitney U test, as appropriate. Differences in categorical variables were analyzed by the chi-square test. Differences in PF volumes in patients with or without LVDD were analyzed by the unpaired Student's *t*-test. Pearson correlations and stepwise multivariate regression analysis among PF volume, metabolic risk factors, and echocardiographic parameters associated with LVDD were performed. Univariate and multivariate backward logistic regression analyses were used to assess the relationships among the presence of LVDD, PF volume, and other risk factors. We also tested the significant association between PF volumes and LVDD in multivariate logistic regression analyses using forced inclusion models of the following parameter: model -1, age and gender; model -2, age, gender, hypertension, and diabetes; model -3, age, gender, hypertension, diabetes, and abdominal obesity. The Hosmer-Lemeshow statistic was applied to assess model calibration. A *P*-value of  $<0.05$  denoted statistical significance, and all tests were two-tailed. Variables were log-transformed if they had a skewed distribution. All analyses were performed using SPSS 17.0J for Windows (SPSS Inc., Tokyo).

195

## Results

### 200 Study sample characteristics

Table 1 shows the characteristics of the participating patients (n=229). Patients with LVDD (n=141) had a higher age, body mass index, waist circumference, systolic  
205 blood pressure, fasting plasma glucose, and hemoglobin A1c, lower estimated glomerular filtration rate; were more likely to have hypertension, diabetes, metabolic syndrome, and coronary artery disease; and used aspirin more frequently than those without LVDD (n=88). In the echocardiographic findings, patients with LVDD had a higher left ventricular mass index and left atrial diameter.

210

### Correlation between PF volume, risk factors, and LVDD

215 PF volume correlated positively with body mass index ( $r=0.51$ ,  $p<0.001$ ), waist circumferences ( $r=0.56$ ,  $p<0.001$ ), systolic blood pressure ( $r=0.16$ ,  $p=0.02$ ), log hemoglobin A1c ( $r=0.23$ ,  $p<0.001$ ), and log triglyceride ( $r=0.32$ ,  $p<0.001$ ), and negatively with high-density lipoprotein cholesterol ( $r=-0.32$ ,  $p<0.001$ ). Multiple stepwise regression analysis identified age ( $\beta=0.13$ ,  $p=0.02$ ) and waist

220 circumference (beta= 0.56,  $p<0.001$ ) as significant and independent correlates with PF  
volume. PF volume correlated significantly with E/e' ( $r=0.21$ ,  $p<0.01$ ; Figure 2-A).  
PF volume also correlated positively with left ventricular mass index ( $r=0.23$ ,  
 $p<0.001$ ; Figure 2-B) and left atrial diameter ( $r=0.32$ ,  $p<0.001$ ; Figure 2-C),  
negatively with the log early/ late diastolic peak flow velocity ratio ( $r=-0.14$ ,  $p=0.03$ ),  
225 but not with LVEF ( $r=-0.07$ ,  $p=0.31$ ). PF volume was larger in patients with LVDD  
( $n=141$ ;  $184\pm 61$  cm<sup>3</sup>) than those without LVDD ( $n=88$ ;  $154\pm 58$  cm<sup>3</sup>,  $p<0.001$ ; Figure  
2-D).

### 230 **Factors associated with LVDD**

Univariate logistic regression analysis found significant relationships between the  
presence of LVDD and age, hypertension, diabetes, estimated glomerular filtration  
235 rate, left ventricular mass index, and PF volume (Table 2). Multivariate backward  
logistic regression analysis identified age (odds ratio [OR]: 1.50, 95% confidence  
interval [CI]: 1.16-1.95 per 10 years,  $p=0.02$ ), the presence of diabetes (OR: 2.24,  
95% CI: 1.16-4.33,  $p=0.02$ ), and PF volume (OR: 2.06 per 100cm<sup>3</sup>, 95% CI:  
1.22-3.47,  $p<0.01$ ) as significant and independent factors associated with the presence  
240 of LVDD. This model was reliable ( $p=0.17$  by the Hosmer-Lemeshow test). In the  
forced entry models, we confirmed the significant association between PF volume and  
the presence of LVDD (OR: 2.09 per 100 cm<sup>3</sup>, 95% CI: 1.15-3.79,  $p= 0.02$ )

independent of age, gender, hypertension, diabetes, and abdominal obesity (Table 3).

This model was also reliable ( $p=0.71$  by the Hosmer-Lemeshow test).

245

250

## Discussion

255

The present study demonstrates that PF volumes correlate significantly with E/e', indicating the association between PF volumes and LVDD. Furthermore, multivariate analysis indicated that PF volume correlates significantly with the presence of LVDD, independent of other risk factors such as age, gender, hypertension, diabetes, or abdominal obesity.

260

PF has been reported to correlate with left atrial dimensions<sup>12</sup> and left ventricular diastolic filling<sup>21</sup>, neither of which are recommended as first-line diagnostic parameters for LVDD<sup>22</sup>. However, recent guidelines recommend the evaluation of tissue Doppler E/e' in the assessment of LVDD<sup>22</sup>. To our knowledge, the present study is the first to report the significant association between PF volume and LVDD using tissue Doppler echocardiography.

265

270

### Potential mechanisms

The potential mechanisms underlying the direct correlation between PF volumes and LVDD are mechanical and paracrine processes. Compression of PF on the

275



myocardium may induce LVDD<sup>12</sup> by disturbing the dilation of the left ventricle and impairing cardiac filling through a pathophysiology similar to the thickening of the pericardium in constrictive pericarditis<sup>23</sup>. Previous studies also reported the paracrine effects of PF. Pericardial and perivascular fat, but not subcutaneous fat, contain high levels of various cytokines<sup>24</sup> that may induce inflammation and increase subsequent collagen turnover, leading to LVDD<sup>25</sup>. Myocardial dysfunction may be induced by a loss of adiponectin secretion or reduced nitric oxide synthase activity in perivascular fat<sup>26</sup>. Both may act directly on cardiomyocytes<sup>27</sup> and indirectly through impaired microvascular relaxation in the myocardium<sup>28</sup>. Thus, it is possible that a similar mechanism operates through paracrine processes. The systemic effects of obesity, diabetes, and other metabolic disorders on LVDD have also been described in previous studies<sup>5</sup>.

Pericardial fat has been observed inside and outside the pericardial sac, which was also reported to be associated with aortic calcification<sup>14</sup>. The pericardial fat outside the sac potentially has some effects on left ventricular function. We measured PF volume by trimming along the pericardial sac using axial, coronal, and sagittal slices and volume-rendered images by 64-slice CT. In the present study, we could not precisely distinguish PF distribution inside or outside the sac. Further studies might be required to determine the effects of the pericardial fat outside the sac on LVDD.

## Cardiac CT and echocardiography

300

We originally quantified PF volume three-dimensionally using multislice CT. This technique yielded highly reproducible measurements (CV = 7.1% and 8.9% for inter- and intra-observer variability, respectively)<sup>7, 29</sup>. In addition, 64-slice CT is commonly  
305 used in cardiovascular clinical practice in Japan.

Noninvasive diagnostic evidence for LVDD is preferably derived from myocardial tissue Doppler. E/e' is reported to correlate closely with LV filling  
310 pressure, and this correlation has been confirmed in patients with pseudo-normal mitral valve flow velocity filling patterns<sup>22</sup>. We defined LVDD simply as E/e' >10 according to the previously described cut-off value<sup>16, 17</sup> partially because of lack of data concerning about duration of reverse pulmonary vein atrial systole flow, duration of mitral valve atrial wave flow, and left atrium volume index used in recent  
315 guideline<sup>22</sup>.

320

## Limitations

325 The clinical study design was cross-sectional, and the results do not imply causality. Because the subjects were not randomly selected but rather inpatients suspected to have coronary artery disease, they were more likely to have coronary artery disease than the general population, their risk factors were considered to have been modified by medications, and some selection bias may influence the result or the present study

330 (B-type natriuretic peptide level and left ventricle mass index are less than expected). The lack of data regarding the extent of coronary artery disease severity is also a limitation because it may largely affect LVDD. E/e' is a marker of increased left ventricular filling pressure, but may occur due to reasons other than primary myocardial disease, affecting the findings. The value of E/e' depends on the

335 technician's experience. Because it depends also on fluid status and medications, the lack of data about time frame between CT and echocardiography may also be limitation. The lack of data about duration of reverse pulmonary vein atrial systole flow, duration of mitral valve atrial wave flow, and left atrial volume index, which are used in recent guideline to diagnose LVDD, is also major limitation on defining

340 LVDD. We have no data about the distribution of body fat (especially abdominal visceral fat and fat outside the pericardial sac) but only have waist circumference though this would help to get more insight into the differential pathophysiologic mechanisms. The correlations between PF volume and parameters of LVDD ( $r = 0.21$  to  $0.32$ ) are weak, although statistically significant, so clinical significance of these

345 correlations should be considered cautiously. The multivariate statistical models  
incorporated both abdominal obesity and PF volume, whose close correlation may  
affect the results of the analyses.

### 350 **Clinical Implications**

In settings where the causality between PF volume and LVDD is demonstrated by a  
prospective study, PF volume may serve as a potential therapeutic target to prevent  
355 LVDD or heart failure in patients with normal ejection fraction (EF). PF volume  
measurement may be useful in selecting therapeutic strategies. For example,  
thiazolidinediones have been reported to increase PF <sup>30</sup> and such medications may not  
be beneficial in patients with LVDD. It may also be useful to assess the effect of  
low-calorie diet on PF volume <sup>31</sup> or the effect of surgical removal <sup>32</sup> of PF on LVDD.

360

### **Conclusions**

365 The present study demonstrated that PF volume correlated significantly with LVDD,  
independent of other risk factors such as age, gender, diabetes, hypertension, or  
abdominal obesity. PF may play a role in the pathogenesis of LVDD.

## **Acknowledgments**

370

The help of Masahiro Hatemura and Kie Shimizu in data analysis is greatly appreciated.

## References

375

1. Tsutsui H, Tsuchihashi-Makaya M, Kinugawa S. Clinical characteristics and outcomes of heart failure with preserved ejection fraction: lessons from epidemiological studies. *J Cardiol.* 2009;55(1):13-22.
- 380 2. Bhatia RS, Tu JV, Lee DS, Austin PC, Fang J, Haouzi A, Gong Y, Liu PP. Outcome of heart failure with preserved ejection fraction in a population-based study. *N Engl J Med.* 2006;355(3):260-269.
3. Grewal J, McCully RB, Kane GC, Lam C, Pellikka PA. Left ventricular function and exercise capacity. *JAMA.* 2009;301(3):286-294.
- 385 4. Jessup M, Brozena S. Heart Failure. *N Engl J Med.* 2003;348(20):2007-2018.
5. Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, Jessup M, Konstam MA, Mancini DM, Michl K, Oates JA, Rahko PS, Silver MA, Stevenson LW, Yancy CW. 2009 Focused update incorporated into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines Developed in Collaboration With the International Society for Heart and Lung Transplantation. *J Am Coll Cardiol.* 2009;53(15):e1-e90.
- 390 6. Kershaw EE, Flier JS. Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab.* 2004;89(6):2548-2556.
- 395 7. Konishi M, Sugiyama S, Sugamura K, Nozaki T, Ohba K, Matsubara J, Matsuzawa Y, Sumida H, Nagayoshi Y, Nakaura T, Awai K, Yamashita Y, Jinnouchi H, Matsui K, Kimura K, Umemura S, Ogawa H. Association of pericardial fat accumulation rather than abdominal obesity with coronary atherosclerotic plaque formation in patients with suspected coronary artery disease. *Atherosclerosis.* 2009;209(2):573-578.
- 400 8. Taguchi R, Takasu J, Itani Y, Yamamoto R, Yokoyama K, Watanabe S, Masuda Y. Pericardial fat accumulation in men as a risk factor for coronary artery disease. *Atherosclerosis.* 2001;157(1):203-209.
- 405 9. Ding J, Hsu FC, Harris TB, Liu Y, Kritchevsky SB, Szklo M, Ouyang P, Espeland MA, Lohman KK, Criqui MH, Allison M, Bluemke DA, Carr JJ. The association of pericardial fat with incident coronary heart disease: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr.* 2009;90(3):499-504.
- 410 10. Katayama T, Fujiwara N, Tsuruya Y. Factors contributing to left atrial enlargement in adults with normal left ventricular systolic function. *J Cardiol.* 2009;55(2):196-204.
11. Liu J, Fox CS, Hickson DA, May WL, Ding J, Carr JJ, Taylor HA. Pericardial fat and echocardiographic measures of cardiac abnormalities: the Jackson Heart Study. *Diabetes Care.* 2011; 34(2):341-346.
- 415 12. Fox CS, Gona P, Hoffmann U, Porter SA, Salton CJ, Massaro JM, Levy D,

- Larson MG, D'Agostino RB, Sr., O'Donnell CJ, Manning WJ. Pericardial fat, intrathoracic fat, and measures of left ventricular structure and function: the Framingham Heart Study. *Circulation*. 2009;119(12):1586-1591.
- 420 **13.** Awai K, Hiraishi K, Hori S. Effect of contrast material injection duration and rate on aortic peak time and peak enhancement at dynamic CT involving injection protocol with dose tailored to patient weight. *Radiology*. 2004;230(1):142-150.
- 425 **14.** Rosito GA, Massaro JM, Hoffmann U, Ruberg FL, Mahabadi AA, Vasan RS, O'Donnell CJ, Fox CS. Pericardial fat, visceral abdominal fat, cardiovascular disease risk factors, and vascular calcification in a community-based sample: the Framingham Heart Study. *Circulation*. 2008;117(5):605-613.
- 15.** Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise J, Solomon S, Spencer KT, St John Sutton M, Stewart W. Recommendations for chamber quantification. *Eur J Echocardiogr*. 2006;7(2):79-108.
- 430 **16.** Redfield MM, Jacobsen SJ, Burnett JC, Jr., Mahoney DW, Bailey KR, Rodeheffer RJ. Burden of systolic and diastolic ventricular dysfunction in the community: appreciating the scope of the heart failure epidemic. *JAMA*. 2003;289(2):194-202.
- 435 **17.** Nagueh SF, Middleton KJ, Kopelen HA, Zoghbi WA, Quinones MA. Doppler tissue imaging: a noninvasive technique for evaluation of left ventricular relaxation and estimation of filling pressures. *J Am Coll Cardiol*. 1997;30(6):1527-1533.
- 440 **18.** Ejima K, Shoda M, Arai K, Suzuki A, Yagishita D, Yagishita Y, Yashiro B, Sato T, Manaka T, Ashihara K, Hagiwara N. Impact of diastolic dysfunction on the outcome of catheter ablation in patients with atrial fibrillation. *Int J Cardiol*. 2011 July 6 [Epub ahead of print].
- 19.** New criteria for 'obesity disease' in Japan. *Circ J*. 2002;66(11):987-992.
- 445 **20.** Imai E, Matsuo S, Makino H, Watanabe T, Akizawa T, Nitta K, Iimuro S, Ohashi Y, Hishida A. Chronic Kidney Disease Japan Cohort (CKD-JAC) study: design and methods. *Hypertens Res*. 2008;31(6):1101-1107.
- 21.** Iacobellis G, Leonetti F, Singh N, A MS. Relationship of epicardial adipose tissue with atrial dimensions and diastolic function in morbidly obese subjects. *Int J Cardiol*. 2007;115(2):272-273.
- 450 **22.** Paulus WJ, Tschope C, Sanderson JE, Rusconi C, Flachskampf FA, Rademakers FE, Marino P, Smiseth OA, De Keulenaer G, Leite-Moreira AF, Borbely A, Edes I, Handoko ML, Heymans S, Pezzali N, Pieske B, Dickstein K, Fraser AG, Brutsaert DL. How to diagnose diastolic heart failure: a consensus statement on the diagnosis of heart failure with normal left ventricular ejection fraction by the Heart Failure and Echocardiography Associations of the European Society of Cardiology. *Eur Heart J*. 2007;28(20):2539-2550.
- 455 **23.** Little WC, Freeman GL. Pericardial Disease. *Circulation*. 2006;113(12):1622-1632.
- 460 **24.** Mazurek T, Zhang L, Zalewski A, Mannion JD, Diehl JT, Arafat H, Sarov-Blat L, O'Brien S, Keiper EA, Johnson AG, Martin J, Goldstein BJ, Shi Y. Human epicardial adipose tissue is a source of inflammatory mediators. *Circulation*.

2003;108(20):2460-2466.

- 465 **25.** Mak GJ, Ledwidge MT, Watson CJ, Phelan DM, Dawkins IR, Murphy NF, Patle AK, Baugh JA, McDonald KM. Natural history of markers of collagen turnover in patients with early diastolic dysfunction and impact of eplerenone. *J Am Coll Cardiol.* 2009;54(18):1674-1682.
- 470 **26.** Greenstein AS, Khavandi K, Withers SB, Sonoyama K, Clancy O, Jeziorska M, Laing I, Yates AP, Pemberton PW, Malik RA, Heagerty AM. Local inflammation and hypoxia abolish the protective anticontractile properties of perivascular fat in obese patients. *Circulation.* 2009;119(12):1661-1670.
- 475 **27.** Silberman GA, Fan TH, Liu H, Jiao Z, Xiao HD, Lovelock JD, Boulden BM, Widder J, Fredd S, Bernstein KE, Wolska BM, Dikalov S, Harrison DG, Dudley SC, Jr. Uncoupled cardiac nitric oxide synthase mediates diastolic dysfunction. *Circulation.* 121(4):519-528.
- 28.** Lohn M, Dubrovskaja G, Lauterbach B, Luft FC, Gollasch M, Sharma AM. Periadventitial fat releases a vascular relaxing factor. *FASEB J.* 2002;16(9):1057-1063.
- 480 **29.** Gorter PM, van Lindert AS, de Vos AM, Meijs MF, van der Graaf Y, Doevendans PA, Prokop M, Visseren FL. Quantification of epicardial and peri-coronary fat using cardiac computed tomography; reproducibility and relation with obesity and metabolic syndrome in patients suspected of coronary artery disease. *Atherosclerosis.* 2008;197(2):896-903.
- 485 **30.** Jonker JT, Lamb HJ, van der Meer RW, Rijzewijk LJ, Menting LJ, Diamant M, Bax JJ, de Roos A, Romijn JA, Smit JW. Pioglitazone compared with metformin increases pericardial fat volume in patients with type 2 diabetes mellitus. *J Clin Endocrinol Metab.* 2009;95(1):456-460.
- 490 **31.** Iacobellis G, Singh N, Wharton S, Sharma AM. Substantial changes in epicardial fat thickness after weight loss in severely obese subjects. *Obesity (Silver Spring).* 2008;16(7):1693-1697.
- 32.** White CM, Sander S, Coleman CI, Gallagher R, Takata H, Humphrey C, Henyan N, Gillespie EL, Kluger J. Impact of epicardial anterior fat pad retention on postcardiothoracic surgery atrial fibrillation incidence: the AFIST-III Study. *J Am Coll Cardiol.* 2007;49(3):298-303.
- 495



## Figure Legends

500 **Figure 1. Representative computed tomography (CT) images of a 55-year-old male patient with hypertension, dyslipidemia, and diabetes.**

Red area is the pericardial fat (PF) within the pericardial sac quantified  
505 three-dimensionally by CT. The axial (A), sagittal (B), and coronal (C) slice of the PF. His PF volume was 265 cm<sup>3</sup> and E/e' was 15.

**Figure 2. Correlation between pericardial fat (PF) volume and left ventricular  
510 diastolic dysfunction (LVDD).**

Correlation between PF volume and echocardiographic parameters: E/e' (A), left  
ventricular mass index (B), and left atrial diameter (C). PF volume had a significant  
515 positive correlation with E/e' (r=0.21, p<0.01), left ventricular mass index (r=0.23,  
p<0.001), and left atrial diameter (r=0.32, p<0.001). Filled and open circles indicate  
patients with and without LVDD, respectively. (D) PF volume in patients with or  
without LVDD. Bar graph indicates mean +standard deviation. PF volume was

significantly larger in patients with LVDD ( $184 \pm 61 \text{ cm}^3$ ) than in those without LVDD  
520 ( $154 \pm 58 \text{ cm}^3$ ,  $p < 0.001$ ).