

Article

Mn(III)-Initiated Facile Oxygenation of Heterocyclic 1,3-Dicarbonyl Compounds

Md. Taifur Rahman, Md. Aminul Haque, Hikaru Igarashi and Hiroshi Nishino *

Department of Chemistry, Graduate School of Science and Technology, Kumamoto University, Kurokami 2-39-1, Kumamoto 860-8555, Japan

* Author to whom correspondence should be addressed; E-Mail: nishino@sci.kumamoto-u.ac.jp; Tel.: +81-96-342-3374; Fax: +81-96-342-3374.

Received: 27 September 2011; in revised form: 7 November 2011 / Accepted: 9 November 2011 / Published: 16 November 2011

Abstract: The Mn(III)-initiated aerobic oxidation of heterocyclic 1,3-dicarbonyl compounds, such as 4-alkyl-1,2-diphenylpyrazolidine-3,5-diones, 1,3-dialkylpyrrolidine-2,4-diones, 3-alkyl-1,5-dimethylbarbituric acids, and 3-butyl-4-hydroxy-2-quinolinone gave excellent to good yields of the corresponding hydroperoxides, which were gradually degraded by exposure to the metal initiator after the reaction to afford the corresponding alcohols. The synthesis of 30 heterocyclic 1,3-dicarbonyl compounds, the corresponding hydroperoxides and the 10 alcohols, their characterization, and the limitations of the procedure are described. In addition, the mechanism of the hydroperoxidation and the redox decomposition of the hydroperoxides are discussed.

Keywords: hydroperoxides; peroxy radicals; aerobic oxidation; redox; 2-hydroxy-1,3-dicarbonyl compounds


1. Introduction

Oxygenation of enolates has been considered as an important organic transformation [1-3], because the α -hydroxycarbonyl moiety is used as a starting material or an intermediate for the synthesis of some natural products [4-8]. In particular, the 2-hydroxy-1,3-dicarbonyl moiety can be found in biologically important compounds such as indole alkaloids, e.g., vindorosine and vindoline [9-12], the cyclopentenoid kjellamanianone [13], and tetracycline-type antibiotics such as doxycycline [14,15]. 4-Butyl-1,2-diphenylpyrazolidine-3,5-dione (phenylbutazone), a nonsteroidal drug, is an efficient

reducing cofactor for the peroxidase activity of prostaglandin H synthase [16,17]. Phenylbutazone inhibits the production of lipid mediators causing inflammation but paradoxically performs this *via* the intermediacy of the peroxy radical and hydroperoxide, which may themselves be proinflammatory. 4-Hydroperoxyphenylbutazone shows a significantly stronger cardiodepressive and coronary constricting effects compared to phenylbutazone itself, 4-hydroxyphenylbutazone, and the ring-opened decomposition product of the hydroperoxide [18]. These phenomena could shed light on the significance of the 4-hydroperoxyphenylbutazone regarding the antiinflammatory or other biological activities of phenylbutazone [19,20] and could explain the side effects such as gastric irritation and toxicity associated with phenylbutazone. Although many reagents have been utilized for the introduction of an oxygen functionality at the 2-position of 1,3-dicarbonyl compounds, the hydroxyl functionality could be introduced at the α -position in most of the cases [21-30].

In connection with our current investigation of the $Mn(OAc)_3$ -assisted aerobic oxidation [31-36], we found the double hydroperoxyalkylation of barbituric acids [37], pyrazolidine-3,5-diones (Scheme 1) [38,39], 4-hydroxy-1*H*-quinolin-2-ones [40], and tetronic acid [41]. The reaction could not be stopped at the monohydroperoxyalkylation stage.

Scheme 1. Mn(III)-assisted aerobic oxidation of 1,2-diphenylpyrazolidine-3,5-dione (R = H) [38-39] and phenylbutazone (R = Bu) [42].



Surprisingly, when phenylbutazone was treated under similar reaction conditions in the absence of an alkene, direct hydroperoxidation occurred and the corresponding solid 4-hydroperoxyphenylbutazone was produced quantitatively (Scheme 1) [42]. We foresaw that the reaction could be very useful for the synthesis of hydroperoxides under very mild aerobic oxidation conditions. Therefore, in order to examine the applicability of $Mn(OAc)_3$ -assisted α -hydroperoxidation, biologically important heterocyclic 1,3-dicarbonyl compounds, e.g., 1,3-dialkylpyrrolidine-2,4-diones **3**, 3-alkyl-1,5-dimethylbarbituric acids **5**, and 3-butyl-4-hydroxy-2-quinolinone (**7**) as well as various 4-monoalkyl-substituted pyrazolidine-3,5-diones **1** were subjected to the reaction, and we obtained very similar results, whereby the corresponding hydroperoxides were obtained in excellent to good yields. In this paper, the full results of the earlier work [42], applications and limitations of the Mn(III)-initiated direct α -hydroperoxidation, and the related reactions are described. Specifically, the full characterization of all the starting materials **1**, **5**, and **7**, and all the hydroperoxides published in the reference [42], the full results of the hydroperoxidation of 1,3-dialkylpyrrolidine-2,4-diones **3**, including the characterization of the corresponding hydroperoxides, and the mechanism of the Mn(III)-initiated autoxidation are discussed. In addition, the limitations of the hydroperoxidation are mentioned using the reaction of acyclic amides, cyclic ketones and esters as model, and the synthesis of 2-hydroxy-1,3-dicarbonyl compounds using the reduction of the hydroperoxides is described.

2. Results and Discussion

2.1. Aerobic Oxidation of 4-Alkyl-substituted Pyrazolidine-3,5-diones **1a–h**

4-Alkyl-1,2-diphenylpyrazolidine-3,5-diones **1a–h** were prepared by a simple condensation reaction between the corresponding alkyl malonate and 1,2-diphenylhydrazine in the presence of NaH in boiling chlorobenzene [17]. With the pyrazolidinediones **1a–h** in hand, a mixture of phenylbutazone (**1e**, 1 mmol) and Mn(OAc)₃ (0.1 mmol) in glacial AcOH (30 mL) was stirred at 23 °C under an aerobic atmosphere, and the desired 4-hydroperoxypyrazolidinedione **2e** was produced in quantitative yield (Scheme 2 and Table 1, entry 9). To scrutinize the role of Mn(OAc)₃ in the aerobic oxidation, we carried out some reactions of 4-*i*-propypyrazolidinedione **1d** under different reaction conditions. When the pyrazolidinedione **1d** was stirred in the absence of Mn(OAc)₃, no conversion to the hydroperoxide **2d** was observed (entry 4). Using Cu(OAc)₂ and ammonium cerium(IV) nitrate (CAN) as the catalyst was not effective in the autoxidation (entries 5 and 6). Therefore, it could be inferred that Mn(OAc)₃ was essential for the hydroperoxidation and that the reaction is neither thermal nor it is induced by sunlight or visible light. While carrying out the reaction of **1d** in the presence or absence of light (entries 7 and 8), there is no alteration in the yield of the hydroperoxide **2d**. Thus, the reaction is not light-assisted. Various alkyl groups were used as the substituent at the 4-position of the pyrazolidinedione, but in all cases similar quantitative conversion to the corresponding hydroperoxides was observed. The results are summarized in Table 1.

Scheme 2. Autoxidation of 4-alkyl-substituted 1,2-diphenylpyrazolidine-3,5-diones **1a–h**.




Table 1. Aerobic oxidation of 4-alkyl-substituted pyrazolidinediones **1a–h** in the presence of a metal oxidant ^a.

Entry	1	Oxidant	1 :oxidant ^b	Time/h	Product/yield (%) ^c
1	1a	Mn(OAc) ₃	1:0.1	2	2a (95)
2	1b	Mn(OAc) ₃	1:0.1	2	2b (95)
3	1c	Mn(OAc) ₃	1:0.1	2	2c (99)
4	1d	none	-	3	Recovery of 1d (100)
5	1d	Cu(OAc) ₂	1:0.2	15	Recovery of 1d (94)
6 ^d	1d	CAN ^e	1:0.1	1	2d (40)
7	1d	Mn(OAc) ₃	1:0.1	2	2d (97)

Table 1. Cont.

Entry	1	Oxidant	1:oxidant ^b	Time/h	Product/yield (%) ^c
8 ^f	1d	Mn(OAc) ₃	1:0.1	2	2d (96)
9	1e	Mn(OAc) ₃	1:0.1	2	2e (99)
10	1f	Mn(OAc) ₃	1:0.1	10 min	2f (95)
11	1g	Mn(OAc) ₃	1:0.1	2	2g (97)
12	1h	Mn(OAc) ₃	1:0.1	2	2h (99)


^a The reaction of **1** (1 mmol) was carried out in AcOH at ambient temperature in air; ^b Molar ratio;

^c Isolated yield is based on the pyrazolidinedione **1** used; ^d The reaction was carried out in MeOH at 0 °C; ^e Ammonium cerium(IV) nitrate; ^f The reaction was carried out in the dark.

2.2. Aerobic Oxidation of 3-Alkyl-substituted Pyrrolidine-2,4-diones **3a–p**

We were pleased to confirm that the specific hydroperoxidation succeeded using pyrazolidinediones **1a–h**, so we next applied the reaction to 1,3-dialkyl-substituted pyrrolidine-2,4-diones **3a–p** which are structurally similar to the pyrazolidinediones. The pyrrolidine-2,4-diones **3a–p** were prepared by the Dieckmann condensation of *N*-alkanoyl-*N*-alkylglycinates which were produced by the reaction of α -bromoacetate with alkylamines followed by alkanoylation with the corresponding alkanoyl chloride (see Experimental section) [43–47]. The 2,4-pyrrolidinediones **3a–p** exist as an enol form (3-alkyl-4-hydroxy-3-pyrrolin-2-ones) in an aprotic polar solvent, such as DMSO-*d*₆ [48]. With the 2,4-pyrrolidinediones **3a–p** in hand, we first explored the hydroperoxidation of **3a**. When the reaction was carried out using a stoichiometric amount of Mn(OAc)₃, the oxidant was consumed after 2 h, and the desired hydroperoxide **4a** was obtained in 65% yield (Scheme 3 and Table 2, entry 1).

Scheme 3. Aerobic oxidation of 1,3-dialkyl-substituted pyrrolidine-2,4-diones **3a–p**.



In order to optimize the hydroperoxidation, the reaction was conducted under various reaction conditions, and we finally obtained the hydroperoxide **4a** in 94% yield under the conditions of room temperature reaction in air for 2 h using a catalytic amount of Mn(OAc)₃ (entry 3). To expand the scope of the reaction, the reaction of other 2,4-pyrrolidinediones **3b–p** was conducted under the

optimized reaction conditions and similar hydroperoxides **4b–p** were obtained in excellent yields (entries 4–18). However, the reactions using 1-ethyl-3-methylpyrrolidine-2,4-dione (**3q**) and ethyl 1-benzylpyrrolidine-2,4-dione-3-carboxylate (**3r**) gave intractable mixtures and the corresponding hydroperoxides were not isolated. In addition, the reaction of 1-benzyl-3-phenylpyrrolidine-2,4-dione (**3s**) failed because of the solubility problem. Although the 4-hydroperoxy-pyrrolidinediones **2a–h** were stable at ambient temperature in air, the 3-hydroperoxy-pyrrolidinediones **4a–p** gradually decomposed within 2 or 3 days and even when stored in a refrigerator at $-20\text{ }^{\circ}\text{C}$ for a half year.

Table 2. Aerobic oxidation of 1,3-dialkyl-substituted pyrrolidinediones **3a–p**^a.



Entry	3	3:Mn(OAc)₃ ^b	Time/h	Product/yield (%) ^c
1	3a	1:1	2	4a (65)
2	3a	1:0.1	14	4a (68)
3	3a	1:0.1	2	4a (94)
4	3b	1:0.1	2	4b (95)
5	3c	1:0.1	2	4c (98)
6	3d	1:0.1	2	4d (93)
7	3e	1:0.1	2	4e (96)
8	3f	1:0.1	1.5	4f (91)
9	3g	1:0.1	1.5	4g (90)
10	3h	1:0.1	1.5	4h (95)
11	3i	1:0.1	1.5	4i (90)
12	3j	1:0.1	1.5	4j (96)
13	3k	1:0.1	1.5	4k (99)
14	3l	1:0.1	1.5	4l (99)
15	3m	1:0.1	1.5	4m (98)
16	3n	1:0.1	1.5	4n (95)
17	3o	1:0.1	1.5	4o (97)
18	3p	1:0.1	1.5	4p (98)

^a The reaction of **3** (1 mmol) was carried out in AcOH (25 mL) at ambient temperature in air;

^b Molar ratio; ^c The yield is based on the amount of **3** used.

2.3. Aerobic Oxidation of 3-Alkyl-1,5-dimethylbarbituric Acids **5a–e** and 3-Butyl-4-hydroxy-2-quinolinone (**7**)

We next investigated the reaction of other biologically important heterocyclic 1,3-dicarbonyl compounds, such as 5-monosubstituted barbituric acids **5a–e** and 3-butyl-4-hydroxy-2-quinolinone **7**. The barbituric acids **5a–e** were prepared by Pd-catalyzed reductive alkylation of 1,3-dimethylbarbituric acid with acetone and aldehydes [49]. The quinolinone **7** was prepared by condensation of aniline and diethyl butylmalonate followed by dehydration (see Experimental section) [50–52]. With the barbituric acids **5a–e** and the quinolinone **7** in hand, the aerobic oxidation was carried out under similar reaction conditions, and very similar results were obtained, giving the corresponding hydroperoxides **6a–e** and **8**, respectively, in excellent to good yields (Scheme 4 and Table 3).

Scheme 4. Aerobic oxidation of barbituric acids **5a–e** and 4-hydroxy-2-quinolinone **7**.**Table 3.** Aerobic oxidation of barbituric acids **5a–e** and 4-hydroxy-2-quinolinone **7**^a.


Entry	Amide	Amide:Mn(OAc) ₃ ^b	Time/h	Product/yield (%) ^c
1	5a	1:0.1	4	6a (80)
2	5b	1:0.1	2	6b (90)
3	5c	1:0.1	4	6c (94)
4	5d	1:0.1	4	6d (94)
5	5e	1:0.1	2	6e (88)
6	7	1:0.1	2	8 (57)

^a The reaction of **5** (1 mmol) was carried out in AcOH (30 mL) at ambient temperature in air;

^b Molar ratio; ^c The yield is based on the amount of **5** or **7** used.

2.4. Mechanism for the Formation of Hydroperoxides **2**, **4**, **6**, and **8**

The aerobic oxidation might be explained by a radical chain mechanism. To rationalize our experimental results, we presume the formation of Mn(III)-enolate complex **A** *in situ* undergoing single-electron transfer (SET) to give 1,3-dicarbonyl radical **B** and the reduced Mn(II) (Scheme 5) [31-36]. This is the initiation step of the radical chain reaction. The 1,3-dicarbonyl radical **B** could be trapped by dissolved molecular oxygen in solution to produce the peroxy radical **C** [53,54]. The radical **C** could simply abstract a hydrogen atom from the cyclic amides to give the product hydroperoxides and another molecule of 1,3-dicarbonyl radical **B**, which continues the radical chain reaction. Since the redox potential (E°) of Mn(III)/Mn(II) is 1.54 V, it seems that the Mn(III) acts as an initiator rather than as a catalyst such as Cu(II)/Cu(I) ($E^\circ = 0.123$ V) [55-67].

Scheme 5. Mn(III)-initiated autoxidation of cyclic amides.

Involvement of the peroxy radical as well as the hydroperoxide intermediate in the course of transition metal-catalyzed autoxidation with different 1,3-dicarbonyl compounds has been proposed earlier [58,59]; however, there is only one report on the detection and identification of such a peroxy radical by electron spin resonance (ESR) [60]. To the best of our knowledge, there are no reports on the isolation and characterization of 2-hydroperoxy-1,3-dicarbonyl compounds in the transition metal-mediated autoxidation of the 1,3-dicarbonyl compounds, except for Hasegawa's work [61]. They showed that introduction of the hydroperoxy functionality at the active methylene position of 1,3-dicarbonyl compounds could be accomplished by dye-sensitized photoreaction of singlet oxygen with the enolic 1,3-dicarbonyl compounds [61]. Although CAN also mediated the hydroperoxidation (Table 1, entry 6), the role of CAN could be accounted for by a similar function of Mn(III) during the reaction [62-64].

2.5. Conversion of the Hydroperoxides **2** and **6** into the Alcohols **9** and **10**

When the aerobic oxidation of the pyrazolidinedione **1d** was carried out for a longer reaction period with a catalytic amount of Mn(OAc)₃ in AcOH or for shorter reaction time in EtOH, a substantial amount of hydroxylated product **9d** was produced along with the hydroperoxide **2d** (Scheme 6 and Table 4, entries 2–4). Furthermore, utilizing a stoichiometric amount of CAN as a stronger oxidant than Mn(OAc)₃ also gave a similar result (entry 5). The formation of the alcohol **9d** might be attributed to the degradation of the corresponding hydroperoxide **2d**. As we have mentioned earlier, metal ion-mediated conversion of the 1,3-dicarbonyl compounds into their corresponding 2-hydroxylated derivatives has been well studied [21-30], nevertheless the mechanism associated with this conversion was not well explored and remained more or less ambiguous. In order to scrutinize the production of **9d**, the hydroperoxide **2d** was stirred at 23 °C in AcOH without the presence of any metal catalyst and under sunlight for 23 h. As a result, no conversion of **2d** to **9d** took place and 100% recovery of **2d** was possible after the removal of the solvent (entry 6). Thus, while many alkylhydroperoxides are known to be sensitive to heat [65,66] and light [67,68], the conversion of **2d** into **9d** is not thermal or photochemical. However, in the presence of a catalytic amount of Mn(OAc)₃ (0.01 equiv.) in the same reaction system, 47% of the **2d** was converted into its corresponding alcohol **9d** (entry 7). Carrying out a similar reaction under darkness did not alter the course of the reaction,

which implies that the Mn(III)-catalyzed degradation is not light-assisted. We then carried out the reaction of **2d** under an argon atmosphere with Mn(OAc)₂, and we found that Mn(OAc)₂ was also effective in the degradation (entry 8). With cerium(IV), 47% of the alcohol **9d** was formed from **2d** (entry 9). A similar degradation was also observed in the reaction of the hydroperoxides **2c**, **2g** (entries 10 and 11), the barbituric acid **5b** and the hydroperoxide **6b** (entries 12–17). Therefore, we could draw the conclusion that the transformation of hydroperoxides **2** and **6** into their corresponding alcohols **9** and **10** was due to a similar redox reactions using copper salt which has been extensively studied [69-71]. If the Mn(III) and Ce(IV) would function such as Cu(II), the mechanism of the degradation might be depicted in Scheme 7.

Scheme 6. Conversion of the hydroperoxides into the corresponding alcohols.





Table 4. Aerobic oxidation of amides under various conditions followed by decomposition ^a.

Entry	Amide	Catalyst	1:Catalyst ^b	Solvent	Time/h	Product/yield (%) ^c	Recovery/%
1	1d	Mn(OAc) ₃	1:0.1	AcOH	2	2d (97)	
2	1d	Mn(OAc) ₃	1:0.1	AcOH	12	2d (87)	9d (8)
3	1d	Mn(OAc) ₃	1:0.2	AcOH	27	2d (52)	9d (46)
4	1d	Mn(OAc) ₃	1:0.28	EtOH	3	2d (54)	9d (41)
5	1d	CAN ^d	1:1	MeOH	1	2d (35)	9d (55)
6	2d	none		AcOH	23		100
7	2d	Mn(OAc) ₃	1:0.01	AcOH	23	9d (47)	32
8 ^e	2d	Mn(OAc) ₂	1:1	AcOH	14	9d (25)	66
9	2d	CAN ^d	1:1	MeOH	2	9d (47)	33
10	2c	Mn(OAc) ₃	1:0.01	AcOH	17	9c (45)	48
11	2g	Mn(OAc) ₃	1:0.01	AcOH	17	9g (40)	28
12	5b	Mn(OAc) ₃	1:0.1	AcOH	3	6b (88)	10b (7)
13	5b	Mn(OAc) ₃	1:0.1	AcOH	5	6b (79)	10b (18)
14	5b	CAN ^d	1:1	MeOH	0.5	6b (47)	10b (43)
15	6b	none		AcOH	12		100
16	6b	Mn(OAc) ₃	1:0.1	AcOH	14	10b (14)	88
17	6b	Mn(OAc) ₃	1:0.4	AcOH	23	10b (40)	50

^a The reaction of an amide (1 mmol) was carried out in AcOH (30 mL) at room temperature in air;


^b Molar ratio; ^c Isolated yield is based on the amide used; ^d The reaction with ammonium cerium(IV) nitrate (CAN) was carried out at 0 °C; ^e The reaction was carried out under an argon atmosphere.

Scheme 7. Degradation of hydroperoxides by a redox system [69].

For M, Mn: n = 3; Ce: n = 4

2.6. Application of the Mn(III)-Mediated Oxygenation

In order to apply the hydroperoxidation to acyclic β -oxoamides, cyclic ketones, and esters, the reactions of malonamides **11a,b**, cyclopentanonecarboxylate derivatives **13a,b**, and substituted dimedones **15** were examined (Scheme 8).

Scheme 8. Aerobic oxidation of malonamides, cyclic β -diketones and β -ketoesters.

Surprisingly, the malonamides **11a,b** were inactive under the aerobic conditions and only afforded the corresponding alcohols **12a** and **12b**, respectively, under an oxygen atmosphere (1 atm) instead of air. Because the reaction of **13a,b** in AcOH gave an intractable mixture, the reaction was carried out in MeOH and the corresponding alcohols **14a,b** were isolated (Scheme 8). The dimedone derivatives **15** did not produce any oxygenated products, but **15** were recovered. Therefore, the hydroperoxidation must be characteristic for the cyclic amide derivatives. We also assumed that the corresponding hydroperoxides of **11** and **13** were too unstable in the solution containing Mn(III), and thus readily decomposed to give the corresponding alcohols **12** and **14**, respectively, according to the redox system mentioned above (Scheme 7).

The hydroperoxides **2a–d** and **6b** could be transformed by reduction into their corresponding alcohols **9a–d** and **10b**, respectively. Simple stirring of equimolar amounts of the hydroperoxide and Ph₃P in Et₂O at room temperature gave the corresponding alcohol after a very simple work-up procedure (see Experimental section). The results are summarized in Table 5.

Table 5. Reduction of the hydroperoxides **2a–d** and **6b** using Ph₃P^a.

Entry	ROOH	ROOH:Ph ₃ P	Time/h	Product/Yield (%) ^b
1	2a	1:1	3	9a (96)
2	2b	1:1	3	9b (97)
3	2c	1:1	3	9c (98)
4	2d	1:1	3	9d (97)
5	6b	1:1	3	10b (80)


^a The reaction of the hydroperoxides **2a–d** and **6b** with Ph₃P was carried out in Et₂O in air at room temperature; ^b Isolated yield is based on **2** or **6b** used.

2.7. Structural Determination of the Hydroperoxides and the Hydroxyl Derivatives

Characterization of the hydroperoxides and the alcohols deserves comment. All the obtained hydroperoxides in CH₂Cl₂ showed a positive potassium iodide-starch test. The structural assignment of the hydroperoxides **2**, **4**, **6**, and **8** was based on their ¹H-NMR, ¹³C-NMR, IR spectra as well as their elemental analyses. For example, the 4-hydroperoxy-pyrazolidinedione **2g** showed a singlet at δ 11.21 ppm in the ¹H-NMR spectrum due to OOH group. In the ¹³C-NMR spectrum, the amide carbonyl carbon appeared at δ 167.7 ppm and a quaternary carbon C-4 bearing the OOH group at δ 86.7 ppm. In addition, the elemental analysis and FAB HRMS supported the molecular formula of C₂₂H₁₈N₂O₄. The structure was finally confirmed by X-ray crystallography. A colorless single crystal of **2g** was successfully grown from CH₂Cl₂-benzene of approximate dimensions of 0.25 × 0.50 × 0.10 mm was mounted on a glass fiber. All measurements were made on an imaging plate diffractometer with graphite monochromated Mo-K α radiation. Cell constants and an orientation matrix for data collection corresponded to a primitive triclinic cell with dimensions were obtained as the triclinic space group *P*-1 with cell constants $a = 10.3233$, $b = 10.4273$, $c = 12.9985$ Å, $V = 1283.0$ Å³, and $\alpha = 95.792$, $\beta = 104.504$, $\gamma = 105.643^\circ$. The structure was solved by direct methods and expanded using Fourier techniques (see Supplementary data). The ORTEP drawing of **2g** is shown in Figure 1. The intramolecular hydrogen-bonding in **2g** could be visualized between the terminal hydroperoxy

oxygen and the carbonyl oxygen, O(4)-O(1) (2.705 Å) [32,33,37-40,42]. The other hydroperoxides obtained from the aerobic oxidation showed similar spectroscopic features.

Figure 1. ORTEP drawing of 4-benzyl-4-hydroperoxy-1,2-diphenylpyrazolidine-3,5-dione (**2g**).



In the case of the alcohol derivatives **9**, **10**, **12**, and **14**, the characteristic spectral features of the OH group were observed in the IR and NMR spectra. For example, **9g** showed an absorption band at 3298 cm^{-1} in the IR spectrum corresponding to the OH group and a singlet at δ 4.88 ppm in the $^1\text{H-NMR}$ spectrum due to the OH group. In the $^{13}\text{C-NMR}$ spectrum, the quaternary carbon C-4 bearing the OH group appeared at δ 75.8 ppm and this is significantly different from that having the OOH group (Table 6) [72]. In addition, the elemental analysis of **9g** supported the molecular formula of $\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}_3$. Therefore, it is easy to distinguish the corresponding alcohols from the hydroperoxides.

Table 6. $^{13}\text{C-NMR}$ chemical shifts of the quaternary carbon bearing the OOH and OH group ^a.













Hydroperoxide	$^{13}\text{C-OOH/ppm}$	Alcohol	$^{13}\text{C-OH/ppm}$
2a 	81.6	9a 	71.1
2b 	85.8	9b 	74.7
2c 	85.3	9c 	74.0
2d 	87.6	9d 	76.3

Table 6. Cont.

2g		86.7	9g		75.8
6b		87.2	10b		76.6

^a The ¹³C-NMR spectrum was measured in CDCl₃.

3. Experimental

3.1. Measurements

Melting points were taken using a Yanagimoto micromelting point apparatus and were not corrected. The NMR spectra were recorded using a JNM AL300 or ECX 500 FT-NMR spectrometer at 300 or 500 MHz for ¹H and 75 or 125 MHz for ¹³C, with tetramethylsilane as the internal standard. The chemical shifts are reported in δ values (ppm) and the coupling constants in Hz. The IR spectra were measured in CHCl₃ or KBr using a Shimadzu 8400 FT IR spectrometer and expressed in cm⁻¹. The EI MS spectra were measured by a Shimadzu QP-5050A gas chromatograph-mass spectrometer with the ionizing voltage of 70 eV. The high-resolution mass spectra and the elemental analysis were performed at the Instrumental Analysis Center, Kumamoto University, Kumamoto, Japan.

3.2. Materials

Manganese(II) acetate tetrahydrate, Mn(OAc)₂•4H₂O, was purchased from Wako Pure Chemical Ind., Ltd. Manganese(III) acetate dihydrate, Mn(OAc)₃•2H₂O, was prepared according to the method described in the literature [73,74]. 4-Alkyl-1,2-diphenylpyrazolidine-3,5-diones **1a–h** were prepared by the condensation of a suitable alkylmalonate with 1,2-diphenylhydrazine in the presence of NaH in boiling chlorobenzene [17]. Methyl 2-oxotetrahydrofuran-3-carboxylate (**13a**) and ethyl 2-oxocyclopentanecarboxylate (**13b**) were purchased from Tokyo Chemical Industry Co., Ltd., and used as received. The physical data for the pyrazolidinediones **1a–h** are given in supplementary data.

1,3-Dialkyl-substituted pyrrolidine-2,4-diones **3a–s** were prepared as follows [75–81]. To a mixture of ethyl (benzylamino)acetate (3.22 mL; 17.4 mmol) and triethylamine (4.85 mL; 34.8 mmol) in CHCl₃ (25 mL) was dropwise-added propanoyl chloride (1.65 mL; 19.2 mmol) at 0 °C over 15 min. After stirring for another 1.5 h at room temperature, the mixture was diluted with CHCl₃ to 50 mL and washed with a 5% aqueous AcOH solution (25 mL), water (50 mL), brine (25 mL), dried over anhydrous MgSO₄, and then concentrated to dryness, affording the liquid propanoyl-protected (benzylamino)acetate with sufficient purity for use in the next step.

To a refluxing suspension of NaH (60% dispersion in mineral oil) (500 mg; 11.03 mmol) and tetrahydrofuran (50 mL) in a 300 mL three-necked flask was dropwise-added the prepared propanoyl-protected (benzylamino)acetate (2.5 g; 10.03 mmol) in tetrahydrofuran (50 mL). After this addition, the mixture was continuously heated under reflux for 12 h. A pale yellow solid was formed during the heating and then filtered under suction. The obtained solid was dissolved in a minimum volume of water and very carefully acidified with 2 M H₂SO₄, giving 1-benzyl-3-methyl-2,4-pyrrolidinedione

(keto form of **3a**) as a crude precipitate. The precipitate was purified by silica gel column chromatography, eluting with a mixture of EtOAc and hexane followed by recrystallization using the appropriate solvent. The other 2,4-pyrrolidinediones **3b–s** were prepared in a manner similar to that described above. The physical data for the pyrrolidinediones **3a–s** are given in the Supplementary Data.

5-Alkyl-1,3-dimethylbarbituric acids **5a–e** were prepared as follows [49]: 1,3-Dimethylbarbituric acid (5 mmol) and 10%Pd-C (0.2 g) were added to a mixture of acetone (20 mL), AcOH (20 mL), and water (10 mL). A few drops of concentrated H₂SO₄ were added and the mixture was heated at 50 °C under hydrogen (2 atm) for 4 h using a glass reactor. The mixture was filtered in a Celite column and the filtrate was extracted with CHCl₃. The extract was washed with water several times and a saturated aqueous solution of NaHCO₃, and then dried over anhydrous MgSO₄. After removal of the solvent, the crude product was recrystallized from Et₂O/hexane, giving 1,3-dimethyl-5-*i*-propylbarbituric acid (**5a**) in 90% yield. A similar reaction of the barbituric acid (5 mmol) with 2-naphthaldehyde (5 mmol) was carried out in MeOH (30 mL) at the reflux temperature for 10 h. After the same work-up, 5-(2-naphthyl)methyl-1,3-dimethylbarbituric acid **5e** was obtained in 88% yield. Other benzyl-substituted barbituric acids **5b–d** were prepared in the same manner. The physical data for the barbituric acids **5a–e** are given in supplementary data.

3-Butyl-4-hydroxy-2-quinolinone **7** was prepared according to the standard procedure described in the literature [50–52]. Aniline (2.80 g; 30 mmol) and diethylbutyl malonate (6.51 g; 30 mmol) were heated at 150 °C for 20 h. Water (30 mL) was added and the mixture was extracted with CH₂Cl₂. The extract was washed with 2 M HCl, water and dried over anhydrous Na₂SO₄. After removal of the solvent, ethyl 2-(phenylcarbamoyl)hexanoate (28 mmol) was obtained. The hexanoate in EtOH (10 mL) was hydrolyzed with 6 M NaOH (20 mL) in EtOH (20 mL) at room temperature for 1 h. After acidifying the solution with 2 M HCl, the precipitates were filtered with suction. The obtained solid acid was dehydrated in PPA, which was poured into water and then neutralized using 6 M NaOH to pH 4, giving **7** as a precipitate (3.67 g; 56%). The physical data for the quinolinone **7** are given in the Supplementary Data.

2-Benzyl-*N*¹,*N*³-dipropylmalonamide (**11a**) and 2-benzyl-*N*¹,*N*³-dibutylmalonamide (**11b**) were prepared by the reaction of diethyl 2-benzylmalonamide (2.50 g) with propylamine and butylamine (50 mL), respectively, at reflux temperature for 24 h followed by removal of the amine unchanged under reduced pressure to give a colorless solid which was recrystallized with CHCl₃/hexane [82]. The physical data for the malonamides **11a** and **11b** are given in the Supplementary Data.

3.3. Mn(III)-Initiated Aerobic Oxidation of Heterocyclic 1,3-Dicarbonyl Compounds **1a–h**, **3a–p**, **5a–e**, and **7**

A typical procedure is as follows: to a solution of the cyclic amide (1 mmol) in glacial AcOH (25 mL), Mn(OAc)₃•2H₂O (26.8 mg; 0.1 mmol) was added. The mixture was stirred at room temperature in air for 1.5–2 h, and then the reaction was quenched by adding water (25 mL) to the mixture. The aqueous reaction mixture was extracted three times with CH₂Cl₂ (30 mL) and the combined extract was washed with water, then a saturated aqueous solution of NaHCO₃, dried over anhydrous MgSO₄, and concentrated to dryness. Although the product was almost pure, it was further purified by silica gel flash column

chromatography, eluting with Et₂O/hexane (7:3 v/v) or EtOAc/hexane (8:2 v/v) if needed. The physical data for the hydroperoxides **2a–h**, **4a–p**, **6a–e**, and **8** are given in the Supplementary Data.

3.4. Conversion of the Hydroperoxides **2** and **6** into the Alcohols **9** and **10**

The redox reaction of the hydroperoxide **2c**, **2d**, **2g**, **6b** (1 mmol) was carried out in AcOH (30 mL) at room temperature in air. The molar ratio of the catalyst and the reaction times are described in Table 4. After the usual work-up, the corresponding alcohols **9c**, **9d**, **9g**, and **10b** were obtained. On the other hand, the reduction of the hydroperoxides **2a–d** and **6b** using Ph₃P is as follows: to the hydroperoxide (0.5 mmol) in Et₂O (10 mL), a solution of Ph₃P (0.5 mmol) in Et₂O (10 mL) was dropwise added and stirred for 3 h at room temperature. Water (20 mL) was added to the reaction mixture and then extracted with CH₂Cl₂. The extract was dried over anhydrous Na₂SO₄ and then evaporated to dryness. Simple recrystallization from Et₂O/hexane gave the pure crystalline alcohol. The physical data for the alcohols **9a–d**, **9g**, and **10b** are given in the Supplementary Data.

3.5. Aerobic Oxidation of **1d**, **2d**, and **5b** Using Ammonium Cerium (IV) Nitrate (CAN)

A typical procedure is as follows: to an ice-cooled solution of the substrate **1d**, **2d**, or **5b** (1 mmol) in MeOH (20 mL), a solution of ammonium cerium(IV) nitrate (CAN) (1 mmol) in MeOH (10 mL) was dropwise-added and stirred in air until the orange color of Ce(IV) disappeared. The reaction was quenched by adding water (25 mL). The aqueous reaction mixture was extracted three times with CH₂Cl₂ (30 mL), and the combined extract was washed with water, saturated brine, dried over anhydrous Na₂SO₄, and then concentrated to dryness. The residue was separated by a silica gel column, eluted with EtOAc/hexane (7:3 v/v). The obtained products **2d**, **9d**, **6b**, and **10b** were further purified by recrystallization from the solvent described above.

3.6. Mn(III)-Initiated Aerobic Oxidation of Acyclic Amides **11a,b**, ketones **13b**, and Esters **13b,15**

The oxidation of **11a** and **11b** (0.7 mmol) was carried out at room temperature in glacial AcOH (10 mL) in the presence of Mn(OAc)₃•2H₂O (0.42 mmol) under an oxygen atmosphere (1 atm) for 29 h. After the usual work-up, the corresponding alcohols **12a** and **12b** were obtained in poor yields. The physical data for the alcohols **12a** and **12b** are given in the Supplementary Data.

The aerobic oxidation of cyclopentanonecarboxylate derivatives **13a** and **13b** was as follows. To a solution of the cyclopentanonecarboxylate (1 mmol) in MeOH (10 mL), a mixture of Mn(OAc)₃•2H₂O (0.1 mmol) in MeOH (20 mL) was dropwise-added, and the mixture was stirred at room temperature for 12 h in air. The reaction mixture was then filtered through the silica gel column (eluted with Et₂O) to remove the catalyst. After removal of all volatile components using an evaporator, the corresponding alcohols **14a** and **14b** were obtained as oil. The physical data for the alcohols **14a** and **14b** are given in the Supplementary Data.

2-Alkyl-5,5-dimethylcyclohexane-1,3-diones **15** (R = Et, Pr, *i*-Pr, Bu, Bn) were prepared by the reaction of dimedone with the corresponding alkylbromide in the presence of sodium ethoxide in boiling ethanol [83]. The Mn(III)-catalyzed aerobic oxidation of **15** was carried out under various conditions; however, no reaction occurred.

4. Conclusions

The facile α -oxygenation of cyclic amides was achieved by Mn(III)-initiated aerobic oxidation under very mild conditions. The direct hydroperoxidation was characteristic for the cyclic amides, such as 4-alkyl-1,2-diphenylpyrazolidine-3,5-diones **1a–h**, 3-alkylpyrrolidine-2,4-diones **3a–p**, 3-alkyl-1,5-dimethylbarbituric acids **5a–e**, and 3-butyl-4-hydroxy-2-quinolinone **7**. It was found that the obtained hydroperoxides gradually underwent redox degradation in the presence of Mn(III)/Mn(II) or Ce(IV)/Ce(III) under the conditions to convert into the corresponding alcohols. Acyclic amides and esters, such as malonamides **11a,b** and cyclopentanonecarboxylates **13a,b**, afforded the alcohols **12a,b** and **14a,b**, probably derived from a similar redox degradation of the corresponding hydroperoxides. Cyclic diketones, such as dimedone **15**, was inactive under the aerobic oxidation conditions. The reason why the hydroperoxides were isolated in this reaction is assumed to be that, (1) it is difficult for the nucleophilic attack of the hydroperoxy group to occur at the amide carbonyl because the electrophilicity of the amide carbonyl carbon should be weak, and (2) the hydroperoxy group is stabilized by intramolecular hydrogen-bonding with the carbonyl group. The hydroperoxides are easily transformed into the corresponding alcohols by normal reduction using Ph_3P .

Antimalarial testing of the hydroperoxide **2g** was also performed using *Plasmodium falciparum* FCR-3 strain (ATCC 30932), and a weak antimalarial activity (14% inhibition) was confirmed. In addition, in order to examine an inhibitory effect toward mRNAs in cells, HeLa cells were treated with the hydroperoxide **2g** or **6b**. However, an appreciable effect on the intracellular distribution of mRNAs was not observed (data not shown).

Supplementary Materials

Supplementary materials can be accessed on: <http://www.mdpi.com/1420-3049/16/11/9562/s1>.

Acknowledgements

This research was supported by Grants-in-Aid for Scientific Research (C), No. 15550039, No. 19550046, and No. 22550041, from the Japan s of Science. We gratefully acknowledge Teruo Shinmyozu, Institute for Materials Chemistry and Engineering, Kyushu University, Japan, and Mikio Yasutake, Graduate School of Science and Engineering, Saitama University, Japan, for the measurement of the high resolution FAB mass spectrum and X-ray analysis, and also thank to Yusuke Wataya, Laboratory of Drug Informatics, Faculty of Pharmaceutical Science, Okayama University, Japan, for the antimalarial testing. We deeply appreciate Tokio Tani and Miss Yuko Azuma, Department of Biological Science, Graduate School of Science and Technology, Kumamoto University, Japan, for the mRNAs inhibitory testing.

References and Notes

1. Chen, B.-C.; Zhou, P.; Davis, F.A.; Ciganek, E. α -Hydroxylation of enolates and silyl enol ethers. *Org. React.* **2003**, *62*, 1-356.

- Vedej, E.; Engler, D.A.; Telschow, J.E. Transition-metal peroxide reactions. Synthesis of α -hydroxycarbonyl compounds from enolates. *J. Org. Chem.* **1978**, *43*, 188-196 and references cited therein.
- Abu-Omar, M.M.; Espenson, J.H. Oxidations of cyclic β -diketones catalyzed by methylrhodium trioxide. *Organometallics* **1996**, *15*, 3543-3549.
- Heathcock, C.H.; Mahain, C.; Schlecht, M.F.; Utawanit, T. A synthetic approach to the quassinoids. *J. Org. Chem.* **1984**, *49*, 3264-3274.
- Davis, F.A.; Clark, C.; Kumar, A.; Chen, B.-C. Asymmetric synthesis of the AB ring segments of daunomycin and 4-demethoxydaunomycin. *J. Org. Chem.* **1994**, *59*, 1184-1190.
- Davis, F.A.; Liu, H.; Chen, B.-C.; Zhou, P. Oxidation of 1,3-dicarbonyl compounds using (camphorylsulfonyl)oxaziridines. *Tetrahedron* **1998**, *54*, 10481-10492.
- Zhang, W.; Watanabe, K.; Cai, X.; Jung, M.E.; Tang, Y.; Zhan, J. Identifying the minimal enzymes required for anhydrotetracycline biosynthesis. *J. Am. Chem. Soc.* **2008**, *130*, 6068-6069.
- Giovanelli, E.; Leroux, S.; Moisan, L.; Carreyre, H.; Thuéry, P.; Buisson, D.-A.; Meddour, A.; Coustard, J.-M.; Thibaudeau, S.; Rousseau, B.; *et al.* On the elucidation of the mechanism of *Vinca* alkaloid fluorination in superacidic medium. *Org. Lett.* **2011**, *13*, 4116-4119.
- Büchi, G.; Matsumoto, K.E.; Nishimura, H. Total synthesis of (\pm)-vindorosine. *J. Am. Chem. Soc.* **1971**, *93*, 3299-3301.
- Andriamialisoa, R.Z.; Langlois, N.; Langlois, Y. A new efficient total synthesis of vindorosine and vindoline. *J. Org. Chem.* **1985**, *50*, 961-967.
- Sasaki, Y.; Kato, D.; Boger, D.L. Asymmetric total synthesis of vindorosine, vindoline, and key vinblastine analogues. *J. Am. Chem. Soc.* **2010**, *132*, 13533-13544.
- Rannoux, C.; Roussi, F.; Martin, M.-T.; Guéritte, F. Elaboration of vinblastine hybrids using a reactive in situ generated *N*-carboxyanhydride. *Org. Biomol. Chem.* **2011**, *9*, 4873-4881.
- Zhu, J.; Klunder, J.H.; Zwanenburg, B. Stereospecific total synthesis of (-)-kjellmanianone and a revision of its absolute configuration. *Tetrahedron Lett.* **1994**, *35*, 2787-2790.
- Charest, M.G.; Lerner, C.D.; Brubaker, J.D.; Siegel, D.R.; Myers, A.G. A convergent enantioselective route to structurally diverse 6-deoxytetracycline antibiotics. *Science* **2005**, *308*, 395-398.
- Mark, G.; Charest, M.G.; Siegel, D.R.; Myers, A.G. Synthesis of (-)-tetracycline. *J. Am. Chem. Soc.* **2005**, *127*, 8292-8293.
- Reed, G.A.; Griffin, I.O.; Eling, T.E. Inactivation of prostaglandin H synthase and prostacyclin synthase by phenylbutazone. Requirement for peroxidative metabolism. *Mol. Pharmacol.* **1985**, *27*, 109-114.
- Vennerstorm, J.L.; Holmes, T.J., Jr. Preparation and evaluation of electrophilic derivatives of phenylbutazone as inhibitors of prostaglandin-*H*-synthase. *J. Med. Chem.* **1987**, *30*, 563-567.
- Mentz, V.P.; Schulz, M.; Kluge, R. Chemische darstellung von phenylbutazon-hydroperoxid und prüfung der substanz auf herzwirksamkeit unter *in-vitro*- und *in-vivo*-bedingungen. *Arzneim.-Frosch.* **1987**, *37*, 1229-1232.
- Rechenberg, H.K.V. *Phenylbutazone*, 2nd ed.; Edward Arnold Ltd.: London, UK, 1962.

20. Chauhan, S.M.S.; Srinivas, K.A.; Mohapatra, P.P. Oxidation of phenylbutazone with hydrogen peroxide catalyzed by 5,10,15,20-tetraarylporphyrinoiron(III) chlorides in dichloromethane. *Ind. J. Chem.* **1999**, *31B*, 724-725 and references cited therein.
21. Demir, A.S.; Jeganathan, A. Selective oxidation of α,β -unsaturated ketones at the α' -position. *Synthesis* **1992**, 235-247.
22. Vedejs, E.; Engler, D.A.; Telschow, J.E. Transition-metal peroxide reactions. Synthesis of α -hydroxycarbonyl compounds from enolates. *J. Org. Chem.* **1978**, *43*, 188-196.
23. Hubert, A.J.; Starcher, P.S. The Baeyer-Villiger oxidation of alkyl oxocyclohexanecarboxylates. *J. Chem. Soc. C* **1968**, 2500-2502.
24. Adam, W.; Smerz, A.K. Nickel-catalyzed hydroxylation of 1,3-dicarbonyl compounds by dimethyldioxirane. *Tetrahedron* **1996**, *52*, 5799-5804.
25. Christoffers, J. Novel manganese-catalyzed α -oxidation of cyclic β -keto esters with molecular oxygen. *J. Org. Chem.* **1999**, *64*, 7668-7669.
26. Christoffers, J.; Wenner, T. Cerium-catalyzed α -oxidation of β -dicarbonyl compounds with molecular oxygen. *SynLett* **2002**, 119-121.
27. Christoffers, J.; Werner, T.; Unger, S.; Frey, W. Preparation of acyloins by cerium-catalyzed, direct hydroxylation of β -dicarbonyl compounds with molecular oxygen. *Eur. J. Org. Chem.* **2003**, 425-431.
28. Baucherel, X.; Levoirier, E.; Uziel, J.; Juge, S. Monohydroxylation of cyclic and acyclic β -keto esters with molecular oxygen catalyzed by cobalt(II) chloride in neutral conditions. *Tetrahedron Lett.* **2000**, *41*, 1385-1387.
29. Nair, V.; Nair, L.G.; Mathew, J. Cerium(IV) mediated oxygenation of dialkyl malonates: A novel synthesis of tartronic acid derivatives. *Tetrahedron Lett.* **1998**, *39*, 2801-2804.
30. Watanabe, T.; Ishikawa, T. Mild air-oxidation of 1,3-dicarbonyl compounds with cesium salts: Novel α -hydroxylation accompanied by partial hydrolysis of malonate derivatives. *Tetrahedron Lett.* **1999**, *40*, 7795-7798.
31. Nishino, H. The facile synthesis of dihydrofurans by the oxidation of olefins with tris(2,4-pentanedionato)manganese(III). *Bull. Chem. Soc. Jpn.* **1985**, *58*, 1922-1927.
32. Tategami, S.; Yamada, T.; Nishino, H.; Korp, J.D.; Kurosawa, K. Formation of 1,2-dioxacyclohexanes by the reaction of alkenes with tris(2,4-pentanedionato)manganese(III) or with manganese(III) acetate. *Tetrahedron Lett.* **1990**, *31*, 6371-6374.
33. Nishino, H.; Tategami, S.; Yamada, T.; Korp, J.D.; Kurosawa, K. Formation of 1, 2-dioxanes by the use of tris (2,4-pentanedionato)-manganese (III) or manganese (III) acetate. *Bull. Chem. Soc. Jpn.* **1991**, *64*, 1800-1809.
34. Asahi, K.; Nishino, H. Facile endoperoxypropellane synthesis by manganese(III) acetate-mediated aerobic oxidation. *Eur. J. Org. Chem.* **2008**, 2404-2416.
35. Tsubusaki, T.; Nishino, H. Formation of 1,2-dioxolanes using Mn(III)-based reaction of various arylacetylenes with 2,4-pentanedione and related reaction. *Tetrahedron* **2009**, *65*, 3745-3752.
36. Haque, M.A.; Nishino, H. Synthesis of peroxy lactones using Mn(III)-catalyzed aerobic oxidation. *Heterocycles* **2011**, *83*, 1783-1805.
37. Qian, C.-Y.; Nishino, H.; Kurosawa, K. Manganese(II) acetate-mediated double 2-hydroperoxyalkylations of barbituric acid and its derivatives. *J. Org. Chem.* **1993**, *58*, 4448-4451.

38. Rahman, M.T.; Nishino, H.; Qian, C.-Y. Synthesis of 4,4-bis(2-hydroperoxyalkyl)pyrazolidine-3,5-diones using manganese(III)-catalyzed autoxidation. *Tetrahedron Lett.* **2003**, *44*, 5225-5228.
39. Rahman, M.T.; Nishino, H. Manganese(III)-based oxidation of 1,2-disubstituted pyrrolidine-3,5-diones in the presence of alkenes. *Tetrahedron* **2003**, *59*, 8383-8392.
40. Kumabe, R.; Nishino, H. A unique peroxide formation based on the Mn(III)-catalyzed aerobic oxidation. *Tetrahedron Lett.* **2004**, *45*, 703-706.
41. Haque, M.A.; Nishino, H. Synthesis of peroxy lactones using Mn(III)-catalyzed aerobic oxidation. *Heterocycles* **2011**, *83*, 1783-1805.
42. Rahman, M.T.; Nishino, H. Manganese(III)-catalyzed facile direct hydroperoxidation of some heterocyclic 1,3-dicarbonyl compounds. *Org. Lett.* **2003**, *5*, 2887-2890.
43. Fugger, J.; Tien, J.M.; Hunsberger, I.M. The preparation of substituted hydrazines. I. Alkylhydrazines via alkylsydnone. *J. Am. Chem. Soc.* **1955**, *77*, 1843-1848.
44. King, J.A.; McMillan, F.H. The preparation of some α -benzylamino- β,β -dialkoxypropionic acid derivatives. *J. Am. Chem. Soc.* **1950**, *72*, 1236-1240.
45. Koech, P.K.; Krische, M.J. Catalytic addition of metallo-aldehyde enolates to ketones: A new C-C bond-forming hydrogenation. *Org. Lett.* **2004**, *6*, 691-694.
46. Speziale, A.J.; Jaworski, E.G. *N*-Substituted glycinate and alaninate esters. *J. Org. Chem.* **1960**, *25*, 728-732.
47. Zhu, Y.; Zou, X.; Hu, F.; Yao, C.; Liu, B.; Yang, H. Synthesis and herbicidal evaluation of novel 3-[(α -Hydroxy-substituted)benzylidene]pyrrolidine-2,4-diones. *J. Agric. Food Chem.* **2005**, *53*, 9566-9570.
48. Chowdhury, F.A.; Nishino, H.; Kurosawa, K. Simple route to azabicyclic peroxides from tetramic acid derivatives using manganese(III)-based molecular oxygen trapping reaction. *Heterocycles* **1999**, *51*, 575-591.
49. Jursic, B.S.; Neumann, D.M. Reductive C-alkylation of barbituric acid derivatives with carbonyl compounds in the presence of platinum and palladium catalysts. *Tetrahedron Lett.* **2001**, *42*, 4103-4107.
50. Detsi, A.; Bardakos, V.; Markopoulos, J.; Igglessi-Markopoulou, O. Reactions of 2-methyl-3,1-benzoxazin-4-one with active methylene compounds: A new route to 3-substituted 4-hydroxyquinolin-2(1*H*)-ones. *J. Chem. Soc. Perkin Trans. 1* **1996**, 2909-2913 and references cited therein.
51. McQuaid, L.A.; Smith, E.C.; Lodge, D.; Pralong, E.; Wikel, J.H.; Calligaro, D.O.; O'Malley, P.J. 3-Phenyl-4-hydroxyquinolin-2(1*H*)-ones: Potent and selective antagonists at the strychnine-insensitive glycine site on the *N*-methyl-D-aspartate receptor complex. *J. Med. Chem.* **1992**, *35*, 3423-3425.
52. Klásek, A.; Mrkvicka, V.; Pevec, A.; Kosmrlj, J. Novel tandem hydration/cyclodehydration of α -thiocyanatoketones to 2-oxo-3-thiazolines. Application to thiazolo[5,4-*c*]quinoline-2,4(3*aH*,5*H*)-dione Synthesis. *J. Org. Chem.* **2004**, *69*, 5646-5651.
53. Ohshima, T.; Sodeoka, M.; Shibasaki, M. Manganese(III)-based oxidative free-radical reaction of α -allyl- β -keto ester with molecular oxygen. *Tetrahedron Lett.* **1993**, *34*, 8509-8512.
54. Lamarque, L.; Méou, A.; Brun, P. Mn(OAc)₃-promoted hydroxylation of α -carbomethoxy- γ -lactones by molecular oxygen. *Can. J. Chem.* **2000**, *78*, 128-132.

55. Lide, D.R. *Handbook of Chemistry and Physics*; CRC press: Boca Raton, FL, USA, 1994; pp. 22-23.
56. Ling, K.-Q.; Lee, Y.; Macikenas, D.; Protasiewicz, J.D.; Sayre, L.M. Copper(II)-mediated autoxidation of *tert*-butylresorcinols. *J. Org. Chem.* **2003**, *68*, 1358-1366 and references cited therein.
57. Brederick, H.; Bauer, G. Hydroperoxide cyclischer β -diketone. *Justus Liebigs Ann. Chem.* **1970**, *739*, 117-120.
58. Tona, M.; Guardiola, M.; Fajari, L.; Messeguer, A. A study on the mechanism and scope of the radical-mediated oxidation of arylacetoacetates. *Tetrahedron* **1995**, *51*, 10041-10052.
59. Nair, V.; Nair, L.G.; Mathew, S. Cerium(IV) mediated oxygenation of dialkyl malonates: A novel synthesis of tartronic acid derivatives. *Tetrahedron Lett.* **1998**, *39*, 2801-2804.
60. Neumann, B.; Müller, S.C.; Hauser, M.J.B.; Steinbock, O.; Simoyi, R.H.; Dalal, N.S. Identification and kinetics study of the peroxy malonyl radical in the aerobic oxidation of malonic acid by cerium(IV). *J. Am. Chem. Soc.* **1995**, *117*, 6372-6373.
61. Yoshioka, M.; Nishioka, T.; Hasegawa, T. Dye-sensitized photooxidation of 6-acyl- and 6-carboalkoxybenzocycloalken-5-ones: Reaction of singlet oxygen with enolic 1,3-dicarbonyl compounds. *J. Org. Chem.* **1993**, *58*, 278-281.
62. Christoffers, J.; Wenner, T. Cerium-catalyzed α -oxidation of β -dicarbonyl compounds with molecular oxygen. *SynLett* **2002**, 119-121.
63. Christoffers, J.; Werner, T.; Unger, S.; Frey, W. Preparation of acyloins by cerium-catalyzed, direct hydroxylation of β -dicarbonyl compounds with molecular oxygen. *Eur. J. Org. Chem.* **2003**, 425-431.
64. Ye, J.-H.; Xue, J.; Ling, K.-Q.; Xu, J.-H. Ceric ammonium nitrate (CAN) mediated novel dimerizations of 4-hydroxyquinolin-2(1*H*)-ones: The first example of reactions of oxygen-centered radicals from 1,3-dicarbonyl compounds. *Tetrahedron Lett.* **1999**, *40*, 1365-1368.
65. Hiatt, R.; Irwin, K.C. Homolytic decompositions of hydroperoxides. V. Thermal decompositions. *J. Org. Chem.* **1968**, *33*, 1436-1441.
66. Stannett, V.; Mesrobian, R.B. The kinetics of the decomposition of tertiary hydroperoxides in solvents. *J. Am. Chem. Soc.* **1950**, *72*, 4125-4130.
67. Boyacı, F.G.; Takaç, S.; Özdamar, T.H. Effects of process parameters on the kinetics of the decomposition of 2-isopropyl naphthalene hydroperoxide into 2-naphthol and acetone. *Rev. Chem. Engl.* **2000**, *16*, 249-299.
68. Dannley, R.L.; Jalics, G. The decomposition of silyl hydroperoxides. *J. Org. Chem.* **1965**, *30*, 3848-3851.
69. Richardson, W.H. Metal ion decomposition of hydroperoxides. IV. Kinetics and products of copper salt catalyzed decomposition of *t*-butyl hydroperoxide. *J. Am. Chem. Soc.* **1966**, *88*, 975-979.
70. Sheldon, R.A.; Kochi, J.K. *Metal Catalyzed Oxidations in Organic Chemistry*; Academic Press: New York, NY, USA, 1981.
71. Hirano, K.; Sakaguchi, S.; Ishii, Y. Radical addition of ethers to alkenes under dioxygen catalyzed by *N*-hydroxyphthalimide (NHPI)/Co(OAc)₂. *Tetrahedron Lett.* **2002**, *43*, 3617-3620.

72. Olah, G.A.; Parker, D.G.; Yoneda, N.; Pelizza, F. Oxyfunctionalization of hydrocarbons. 1. Protolytic cleavage-rearrangement reactions of tertiary alkyl hydroperoxides with magic acid. *J. Am. Chem. Soc.* **1976**, *98*, 2245-2250.
73. Andrusis, P.J., Jr.; Dewar, M.J.S.; Dietz, R.; Hunt, R.L. Aromatic oxidation by electron transfer. I. Oxidations of *p*-methoxytoluene. *J. Am. Chem. Soc.* **1966**, *88*, 5473-5478.
74. Heiba, E.I.; Dessau, R.M.; Koehl, W.J., Jr. Oxidation by metal salts. III. Reaction of manganic acetate with aromatic hydrocarbons and the reactivity of the carboxymethyl radical. *J. Am. Chem. Soc.* **1969**, *91*, 138-145.
75. Fugger, J.; Tien, J.M.; Hunsberger, I.M. The preparation of substituted hydrazines. I. Alkylhydrazines via alkylsydnone. *J. Am. Chem. Soc.* **1955**, *77*, 1843-1848.
76. King, J.A.; McMillan, F.H. The preparation of some α -benzylamino- β,β -dialkoxypropionic acid derivatives. *J. Am. Chem. Soc.* **1950**, *72*, 1236-1240.
77. Koech, P.K.; Krische, M.J. Catalytic addition of metallo-aldehyde enolates to ketones: A new C-C bond-forming hydrogenation. *Org. Lett.* **2004**, *6*, 691-694.
78. Speziale, A.J.; Jaworski, E.G. *N*-Substituted glycinate and alaninate esters. *J. Org. Chem.* **1960**, *25*, 728-732.
79. Zhu, Y.; Zou, X.; Hu, F.; Yao, C.; Liu, B.; Yang, H. Synthesis and herbicidal evaluation of novel 3-[(α -hydroxy-substituted)benzylidene]pyrrolidine-2,4-diones. *J. Agric. Food Chem.* **2005**, *53*, 9566-9570.
80. Lee, V.J.; Branfman, A.R.; Herrin, T.R.; Rinehart, K.L. Synthesis of 3-dienoyl tetramic acids related to streptolydigin and tirandamycin. *J. Am. Chem. Soc.* **1978**, *100*, 4225-4236.
81. Mallinger, A.; Nadal, B.; Chopin, N.; Gall, T.L. One-pot synthesis of 3-aryltetramic acids. *Eur. J. Org. Chem.* **2010**, 1142-1148.
82. Nemoto, H.; Kubota, Y.; Yamamoto, Y. Development of a new acyl anion equivalent for the preparation of masked activated esters, and their use to prepare a dipeptide. *J. Org. Chem.* **1990**, *55*, 4515-4516.
83. Martínez, R.; Clara-Sosa, A.; Apan, M.T.R. Synthesis and cytotoxic evaluation of new (4,5,6,7-tetrahydro-indol-1-yl)-3-*R*-propionic acids and propionic acid ethyl esters generated by molecular mimicry. *Bioorg. Med. Chem.* **2007**, *15*, 3912-3918.

Sample Availability: Samples are available from the authors.

© 2011 by the authors; licensee MDPI, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

Supplementary Data

The physical data for the pyrazolidinediones **1a–h** are given below.

4-Methyl-1,2-phenylpyrazolidine-3,5-dione (1a). Colorless needles (from CHCl₃-hexane); mp 111–113 °C; IR (KBr) ν 1751, 1716 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.32–7.12 (10H, m, arom H), 3.37 (1H, q, J = 7.7 Hz, -CH-), 1.55 (3 H, d, J = 7.7 Hz, Me); ¹³C-NMR (75 MHz, CDCl₃) δ 170.7 (C=O), 135.6 (arom C), 128.9, 126.7, 122.4 (arom CH), 41.1 (C-4), 12.0 (Me). Anal. Calcd for C₁₆H₁₄N₂O₂: C, 72.16; H, 5.30; N, 10.52. Found: C, 72.01; H, 5.18; N, 10.39.

4-Ethyl-1,2-diphenylpyrazolidine-3,5-dione (1b). Colorless microcrystals (from CH₂Cl₂-hexane); mp 104 °C (lit, mp 105–106 °C); IR (KBr) ν 1751, 1710 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.35–7.12 (10H, m, arom H), 3.34 (1H, t, J = 5.4 Hz, -CH-), 2.15–2.09 (2H, m, -CH₂-CH₃) 1.06 (3H, t, J = 7.5 Hz, -CH₂-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 170.1 (C=O), 135.6 (arom C), 128.8, 126.6, 122.4 (arom CH), 47.1 (C-4), 21.6 (-CH₂-CH₃, Et), 10.1 (-CH₂-CH₃, Et).

1,2-Diphenyl-4-propylpyrazolidine-3,5-dione (1c). Colorless microcrystals (from CHCl₃-hexane); mp 103 °C; IR (KBr) ν 1753, 1720(C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.31–7.13 (10H, m, arom H), 3.39 (1H, t, J = 6.0 Hz, -CH-), 2.10–2.02 (2H, m, -CH₂-C₂H₅), 1.58–1.50 (2H, m, -CH₂-CH₂-CH₃), 0.98–0.93 (3H, m, -CH₂-CH₂-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 170.4 (C=O), 135.7 (arom C), 128.9, 126.7, 122.5 (arom CH), 46.1 (C-4), 30.1 (-CH₂-CH₂-CH₃), 19.2 (-CH₂-CH₂-CH₃), 13.8 (-CH₂-CH₂-CH₃). Anal. Calcd for C₁₈H₁₈N₂O₂: C, 73.45; H, 6.16; N, 9.52. Found: C, 73.46; H, 6.06; N, 9.43.

*1,2-Diphenyl-4-*i*-propylpyrazolidine-3,5-dione (1d)*. Colorless needles (from Et₂O-hexane); mp 142 °C (lit, mp 144–145 °C); IR (KBr) ν 1749, 1714 (C=O); δ 7.31–7.13 (10H, m, arom H), 3.24 (1H, d, J = 3.6 Hz, -CH-), 2.65–2.58 (1H, m, CH₃-CH-CH₃), 1.19 (6H, d, J = 6.9 Hz, CH₃-CH-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 169.8 (C=O), 135.7 (arom C), 128.9, 126.7, 122.5 (arom CH), 51.8 (C-4), 30.6 (CH₃-CH-CH₃), 19.0 (CH₃-CH-CH₃). Anal. Calcd for C₁₈H₁₈N₂O₂: C, 73.45; H, 6.16; N, 9.52. Found: C, 73.37; H, 6.12; N, 9.49.

4-Butyl-1,2-diphenylpyrazolidine-3,5-dione (1e). Colorless needles; mp 104–106 °C (from EtOH) (lit, mp 105.5–106.5 °C); IR (KBr) ν 1751, 1716 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.35–7.15 (10H, m, arom H), 3.40–3.36 (1H, m, -CH-), 2.12–2.04 (2H, m, -CH₂-C₃H₇), 1.51–1.32 (4H, m, -CH₂-CH₂-CH₂-CH₃), 0.92–0.87 (3H, m, -CH₂-CH₂-CH₂-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 170.3 (C=O), 135.7 (arom C), 128.9, 126.7, 122.5 (arom CH), 46.2 (C-4), 27.9 (-CH₂-CH₂-CH₂-CH₃), 22.4 (-CH₂-CH₂-CH₂-CH₃ and -CH₂-CH₂-CH₂-CH₃), 13.6 (-CH₂-CH₂-CH₂-CH₃).

*4-*t*-Butyl-1,2-diphenylpyrazolidine-3,5-dione (1f)*. Colorless needles (from Et₂O-hexane); mp 178 °C; IR (KBr) ν 1743, 1708 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.32–7.12 (10H, m, arom H), 2.95 (1H, s, -CH-), 1.25 (9H, s, 3 × CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 169.4 (C=O), 135.8 (arom C), 128.8, 126.5, 122.5 (arom CH), 55.1 (C-4), 36.7 (-C(CH₃)₃), 27.5 (-C(CH₃)₃). Anal. Calcd for C₁₉H₂₀N₂O₂: C, 74.00; H, 6.54; N, 9.08. Found: C, 74.16; H, 6.55; N, 9.07.

4-Benzyl-1,2-diphenylpyrazolidine-3,5-dione (1g). Colorless blocks (from CHCl₃-hexane); mp 136–138 °C (lit, mp 135–137 °C); IR (KBr) ν 1747, 1718 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ

7.26–6.86 (15H, m, arom H), 3.65 (1H, t, $J = 4.5$ Hz, $-\underline{\text{CH}}-$), 3.44 (2H, d, $J = 4.5$ Hz, $-\text{CH}_2-$); ^{13}C -NMR (75 MHz, CDCl_3) δ 169.3 (C=O), 135.3, 135.0 (arom C), 129.9, 128.7, 128.6, 127.3, 127.0, 123.2 (arom CH), 48.5 (C-4), 33.9 ($-\underline{\text{CH}}_2-$). Anal. Calcd for $\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}_2$: C, 77.17; H, 5.30; N, 8.18. Found: C, 77.16; H, 5.20; N, 8.08.

4-Cyclopentyl-1,2-diphenylpyrazolidine-3,5-dione (1h): Colorless microcrystals (from CH_2Cl_2 -hexane); mp 170 °C; IR (KBr) ν 1749, 1720 (C=O); ^1H -NMR (300 MHz, CDCl_3) δ 7.30–7.12 (10H, m, arom H), 3.41 (1H, d, $J = 4.5$ Hz, $-\underline{\text{CH}}-$), 2.60–2.59 (1H, m, $-(\text{CH}_2)_2-\underline{\text{CH}}-(\text{CH}_2)_2-$), 1.89–1.55 (8H, m, $-(\underline{\text{CH}}_2)_4-$); ^{13}C -NMR (75 MHz, CDCl_3) δ 170.0 (C=O), 135.7 (arom C), 128.8, 126.6, 122.4 (arom CH), 48.5 (C-4), 40.7 ($-(\text{CH}_2)_2-\underline{\text{CH}}-(\text{CH}_2)_2-$), 28.7, 24.8 ($-(\underline{\text{CH}}_2)_4-$). Anal. Calcd for $\text{C}_{20}\text{H}_{20}\text{N}_2\text{O}_2$: C, 74.98; H, 6.29; N, 8.74. Found: C, 74.94; H, 6.23; N, 8.69.

The physical data for the pyrrolidinediones **3a–s** are given below.

1-Benzyl-4-hydroxy-3-methyl-3-pyrrolin-2-one (3a). $R_f = 0.57$ (EtOAc:hexane = 9:1 v/v); colorless needles (from EtOH/hexane); mp 146–150 °C; IR (CHCl_3) ν 3400–3100 (OH), 1784, 1697 (C=O); ^1H -NMR ($\text{DMSO}-d_6$) δ 10.65 (1H, s, OH), 7.15–7.35 (5H, m, arom H), 4.46 (2H, s, $\text{CH}_2-\text{C}=\text{O}$), 3.63 (2H, s, PhCH_2), 1.58 (3H, s, Me); ^{13}C -NMR ($\text{DMSO}-d_6$) δ 173.2 (C-2, C=O), 164.8 (C-4), 138.4, 128.6, 127.4, 127.1 (arom C), 100.2 (C-3), 49.0 (C-5, CH_2), 44.6 (PhCH_2), 6.4 (Me). FAB HRMS (acetone/NBA) calcd for $\text{C}_{12}\text{H}_{14}\text{NO}_2$ 204.1025 (M+H). Found 204.1002.

1-Benzyl-3-ethyl-4-hydroxy-3-pyrrolin-2-one (3b). $R_f = 0.61$ (EtOAc:hexane = 7:3 v/v); colorless cubes (from EtOH/hexane); mp 117–118 °C; IR (CHCl_3) ν 3300–3000 (weak OH), 1770, 1693 (C=O); ^1H -NMR ($\text{DMSO}-d_6$) δ 10.70 (1H, br s, OH), 7.16–7.36 (5H, m, arom H), 4.46 (2H, s, $\text{CH}_2-\text{C}=\text{O}$), 3.64 (2H, s, PhCH_2), 2.11 (2H, q, $J = 7.8$ Hz, CH_2), 0.98 (3H, t, $J = 7.8$ Hz, Me); ^{13}C -NMR ($\text{DMSO}-d_6$) δ 172.7 (C-2, C=O), 164.3 (C-4), 138.3, 128.5, 127.3, 126.9 (arom C), 106.2 (C-3), 48.8 (C-5, CH_2), 44.5 (PhCH_2), 14.5 (CH_2), 13.0 (Me). Anal. Calcd for $\text{C}_{13}\text{H}_{15}\text{NO}_2 \cdot 1/3 \text{H}_2\text{O}$: C, 69.93; H, 7.07; N, 6.27. Found: C, 69.67; H, 6.86; N, 6.30.

1-Benzyl-4-hydroxy-3-propyl-3-pyrrolin-2-one (3c). $R_f = 0.75$ (EtOAc:hexane = 7:3 v/v); colorless needles (from CHCl_3 /hexane); mp 124 °C; IR (CHCl_3) ν 3300–3100 (OH), 1768, 1689 (C=O); ^1H -NMR ($\text{DMSO}-d_6$) δ 10.65 (1H, br s, OH), 7.16–7.36 (5H, m, arom H), 4.47 (2H, s, $\text{CH}_2-\text{C}=\text{O}$), 3.65 (2H, s, PhCH_2), 2.07 (2H, t, $J = 7.5$ Hz, CH_2), 1.41 (2H, m, CH_2), 0.85 (3H, t, $J = 7.5$ Hz, Me); ^{13}C -NMR ($\text{DMSO}-d_6$) δ 172.9 (C-2, C=O), 164.9 (C-4), 138.4, 128.6, 127.4, 127.1 (arom C), 104.8 (C-3), 48.9 (C-5, CH_2), 44.6 (PhCH_2), 23.3, 21.3 (CH_2), 13.9 (Me). FAB HRMS (acetone/NBA) calcd for $\text{C}_{14}\text{H}_{18}\text{NO}_2$ 232.1338 (M+H). Found 232.1340.

1-Benzyl-3-butyl-4-hydroxy-3-pyrrolin-2-one (3d). $R_f = 0.44$ (EtOAc:hexane = 5:5 v/v); colorless needles (from EtOH/ H_2O); mp 101–103 °C; IR (CHCl_3) ν 3300–3000 (OH), 1770, 1689 (C=O); ^1H -NMR ($\text{DMSO}-d_6$) δ 10.59 (1H, br s, OH), 7.38–7.16 (5H, m, arom H), 4.47 (2H, s, $\text{CH}_2-\text{C}=\text{O}$), 3.65 (2H, s, PhCH_2), 2.11 (2H, t, $J = 4.5$ Hz, CH_2), 1.40 (2H, m, CH_2), 1.28 (2H, m, CH_2), 0.88 (3H, t, $J = 4.5$ Hz, Me); ^{13}C -NMR ($\text{DMSO}-d_6$) δ 172.8 (C-2, C=O), 164.7 (C-4), 138.3, 128.5, 127.1, 126.9

(arom C), 104.8 (C-3), 48.8 (C-5, CH₂), 44.8 (PhCH₂), 30.1, 21.9, 20.8 (CH₂), 13.7 (Me). FAB HRMS (acetone/NBA) calcd for C₁₅H₂₀NO₂ 246.1494 (M+H). Found 246.1487.

l-Benzyl-3-hexyl-4-hydroxy-3-pyrrolin-2-one (**3e**). *R*_f = 0.62 (EtOAc:hexane = 5:5 v/v); colorless needles (from EtOH/hexane); mp 94–95 °C; IR (CHCl₃) ν 3600–3300 (weak OH), 1768, 1689 (C=O); ¹H-NMR (DMSO-*d*₆) δ 10.68 (1H, br s, OH), 7.19–7.39 (5H, m, arom H), 4.50 (2H, s, CH₂-C=), 3.68 (2H, s, PhCH₂), 2.13 (2H, t, *J* = 9.0 Hz, CH₂), 1.29–1.46 (8H, m, 4CH₂), 0.89 (3H, t, *J* = 9.0 Hz, Me); ¹³C-NMR (DMSO-*d*₆) δ 173.0 (C-2, C=O), 164.8 (C-4), 138.4, 128.6, 127.4, 127.1 (arom C), 105.0 (C-3), 48.6 (C-5, CH₂), 44.6 (PhCH₂), 31.2, 28.6, 27.9, 22.2, 21.2 (CH₂), 13.9 (Me). FAB HRMS (acetone/NBA) calcd for C₁₇H₂₄NO₂ 274.1807 (M+H). Found 274.1795.

l-Butyl-4-hydroxy-3-methyl-3-pyrrolin-2-one (**3f**). *R*_f = 0.32 (CHCl₃:MeOH = 98:2 v/v); colorless needles (from ethanol/hexane); mp 110–112 °C; IR (CHCl₃) ν 3250–3000 (weak OH), 1772, 1685 (C=O); ¹H-NMR (DMSO-*d*₆) δ 10.56 (1H, br s, OH), 3.71 (2H, s, CH₂-C=), 3.23 (2H, t, *J* = 6.9 Hz, CH₂), 1.53 (3H, s, Me), 1.43 (2H, m, CH₂), 1.24 (2H, m, CH₂), 0.87 (3H, t, *J* = 7.2 Hz, Me); ¹³C-NMR (DMSO-*d*₆) δ 172.9 (C-2, C=O), 164.2 (C-4), 100.3 (C-3), 49.2 (C-5, CH₂), 44.6 (PhCH₂), 40.3, 30.2, 19.5 (CH₂), 13.7 (Me), 6.4 (Me). FAB HRMS (acetone/NBA) calcd for C₉H₁₆NO₂ 170.1181 (M+H). Found 170.1178.

l-Butyl-3-ethyl-4-hydroxy-3-pyrrolin-2-one (**3g**). *R*_f = 0.49 (EtOAc:hexane = 6:4 v/v); colorless needles (from EtOAc/hexane); mp 85–87 °C; IR (CHCl₃) ν 3500–3100 (OH), 1768, 1685 (C=O); ¹H-NMR (DMSO-*d*₆) δ 10.47 (1H, br s, OH), 3.70 (2H, s, CH₂-C=), 3.23 (2H, t, *J* = 6.9 Hz, CH₂), 2.04 (2H, q, *J* = 7.2 Hz, CH₂), 1.43 (2H, m, CH₂), 1.21 (2H, m, CH₂), 0.89 (6H, t, *J* = 7.2 Hz, 2Me); ¹³C-NMR (DMSO-*d*₆) δ 172.4 (C-2, C=O), 163.8 (C-4), 106.4 (C-3), 48.9 (C-5, CH₂), 40.3, 30.1, 19.4, 14.4 (CH₂), 13.7 (Me), 6.4 (Me). FAB HRMS (acetone/NBA) calcd for C₁₀H₁₈NO₂ 184.1338 (M+H). Found 184.1338.

l,3-Dibutyl-4-hydroxy-3-pyrrolin-2-one (**3h**). *R*_f = 0.69 (EtOAc:hexane = 6:4 v/v); colorless microcrystals (from cold water); mp 113–114 °C; IR (CHCl₃) ν 3300–3100 (OH), 1770, 1693 (C=O); ¹H-NMR (DMSO-*d*₆) δ 10.59 (1H, br s, OH), 3.77 (2H, s, CH₂-C=), 3.29 (2H, t, *J* = 6.9 Hz, CH₂), 2.09 (2H, t, *J* = 7.5 Hz, CH₂), 1.29–1.49 (8H, m, 4CH₂), 0.93 (6H, t, *J* = 7.2 Hz, 2Me); ¹³C-NMR (DMSO-*d*₆) δ 172.8 (C-2, C=O), 164.5 (C-4), 105.3 (C-3), 49.1 (C-5, CH₂), 40.3, 30.1, 21.9, 20.7, 19.4 (CH₂), 13.7 (Me), 13.5 (Me). FAB HRMS (acetone-NBA) calcd for C₁₂H₂₂NO₂ 212.1651 (M+H). Found 212.1635.

l-(*t*-Butyl)-4-hydroxy-3-methyl-3-pyrrolin-2-one (**3i**). *R*_f = 0.72 (EtOAc:hexane = 8:2 v/v); colorless needles (from EtOAc/hexane); mp 184 °C; IR (CHCl₃) ν 3300–3000 (weak OH), 1768, 1689 (C=O); ¹H-NMR (DMSO-*d*₆) δ 10.36 (1H, br s, OH), 3.76 (2H, s, CH₂-C=), 1.48 (3H, s, Me), 1.32 (9H, s, 3Me); ¹³C-NMR (DMSO-*d*₆) δ 173.3 (C-2, C=O), 163.4 (C-4), 101.6 (C-3), 52.3 (quart C), 47.5 (C-5, CH₂), 27.7 (3Me), 6.2 (Me). FAB HRMS (acetone-NBA) calcd for C₉H₁₆NO₂ 170.1181 (M+H). Found 170.1179.

l-(*t*-Butyl)-3-ethyl-4-hydroxy-3-pyrrolin-2-one (**3j**). *R*_f = 0.32 (EtOAc:hexane = 4:6 v/v); colorless needles (from CHCl₃/hexane); mp 129–130 °C; IR (CHCl₃) ν 3300–3100 (weak OH), 1768, 1683

(C=O); $^1\text{H-NMR}$ (DMSO- d_6) δ 10.37 (1H, br s, OH), 3.76 (2H, s, $\text{CH}_2\text{-C=}$), 2.00 (2H, q, $J = 7.5$ Hz, CH_2), 1.32 (9H, s, 3Me), 0.92 (3H, t, $J = 7.5$ Hz, Me); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 172.9 (C-2, C=O), 163.1 (C-4), 107.6 (C-3), 52.3 (quart C), 47.4 (C-5, CH_2), 27.7 (3Me), 14.3 (CH_2), 13.1 (Me). FAB HRMS (acetone/NBA) calcd for $\text{C}_{10}\text{H}_{18}\text{NO}_2$ 184.1338 (M+H). Found 184.1325.

l-(*t*-Butyl)-4-hydroxy-3-propyl-3-pyrrolin-2-one (**3k**). $R_f = 0.64$ (EtOAc:hexane = 5:5 v/v); colorless needles (from EtOH/hexane); mp 145–146 °C; IR (CHCl_3) ν 3400–3100 (weak OH), 1766, 1687 (C=O); $^1\text{H-NMR}$ (DMSO- d_6) δ 10.31 (1H, br s, OH), 3.77 (2H, s, $\text{CH}_2\text{-C=}$), 1.97 (2H, t, $J = 7.8$ Hz, CH_2), 1.38 (2H, m, CH_2), 1.36 (9H, s, 3Me), 0.82 (3H, t, $J = 7.2$ Hz, Me); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 173.1 (C-2, C=O), 163.6 (C-4), 106.2 (C-3), 52.3 (quart C), 47.4 (C-5, CH_2), 27.7 (3Me), 23.1, 21.2 (CH_2), 13.9 (Me). FAB HRMS (acetone/NBA) calcd for $\text{C}_{11}\text{H}_{20}\text{NO}_2$ 198.1494 (M+H). Found 198.1487.

l-(*t*-Butyl)-3-butyl-4-hydroxy-3-pyrrolin-2-one (**3l**). $R_f = 0.59$ (EtOAc:hexane = 5:5 v/v); colorless needles (from EtOH/hexane); mp 142–144 °C; IR (CHCl_3) ν 3500–3050 (OH), 1766, 1687 (C=O); $^1\text{H-NMR}$ (DMSO- d_6) δ 10.33 (1H, br s, OH), 3.76 (2H, s, $\text{CH}_2\text{-C=}$), 1.99 (2H, t, $J = 4.2$ Hz, CH_2), 1.29 (9H, s, 3Me), 1.23 (4H, m, 2 CH_2), 0.82 (3H, t, $J = 7.2$ Hz, Me); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 173.2 (C-2, C=O), 163.6 (C-4), 106.4 (C-3), 52.4 (quart C), 47.5 (C-5, CH_2), 27.8 (3Me), 30.1, 22.1, 20.8 (CH_2), 13.8 (Me). FAB HRMS (acetone/NBA) calcd for $\text{C}_{12}\text{H}_{22}\text{NO}_2$ 212.1651 (M+H). Found 212.1637.

l-(*t*-Butyl)-3-hexyl-4-hydroxy-3-pyrrolin-2-one (**3m**). $R_f = 0.53$ (EtOAc:hexane = 3:7 v/v); colorless needles (from EtOH/hexane); mp 94 °C; IR (CHCl_3) ν 3300–3100 (weak OH), 1780, 1683 (C=O); $^1\text{H-NMR}$ (DMSO- d_6) δ 10.32 (1H, br s, OH), 3.74 (2H, s, $\text{CH}_2\text{-C=}$), 1.95 (2H, t, $J = 7.8$ Hz, CH_2), 1.36 (9H, s, 3Me), 1.36–1.21 (8H, m, 4 CH_2), 0.83 (3H, t, $J = 7.2$ Hz, Me); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 173.2 (C-2, C=O), 163.5 (C-4), 106.4 (C-3), 52.4 (quart C), 47.4 (C-5, CH_2), 31.1, 28.7 (CH_2), 27.9 (3Me), 27.8, 22.1, 21.0 (CH_2), 13.9 (Me). FAB HRMS (acetone/NBA) calcd for $\text{C}_{14}\text{H}_{26}\text{NO}_2$ 240.1964 (M+H). Found 240.1948.

4-Hydroxy-3-methyl-1-(*i*-propyl)-3-pyrrolin-2-one (**3n**). $R_f = 0.30$ (EtOAc:hexane = 7:3 v/v); colorless needles (from EtOAc/hexane); mp 140–142 °C; IR (CHCl_3) ν 3300–3100 (weak OH), 1770, 1683 (C=O); $^1\text{H-NMR}$ (DMSO- d_6) δ 10.56 (1H, br s, OH), 4.15 (1H, sep, $J = 6.9$ Hz, $-\text{CH}<$), 3.66 (2H, s, $\text{CH}_2\text{-C=}$), 1.52 (3H, s, Me), 1.06 (6H, d, $J = 6.9$ Hz, 2Me); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 72.3 (C-2, C=O), 164.4 (C-4), 100.5 (C-3), 44.5 (C-5, CH_2), 41.1 (CH), 20.3 (2Me), 6.2 (Me). FAB HRMS (acetone/NBA) calcd for $\text{C}_8\text{H}_{14}\text{NO}_2$ 156.1025 (M+H). Found 156.1023.

3-Ethyl-4-hydroxy-1-(*i*-propyl)-3-pyrrolin-2-one (**3o**): $R_f = 0.55$ (EtOAc:hexane = 7:3 v/v); colorless needles (from EtOAc/hexane); mp 117–118 °C; IR (CHCl_3) ν 3300–3100 (weak OH), 1770, 1683 (C=O); $^1\text{H-NMR}$ (DMSO- d_6) δ 10.52 (1H, br s, OH), 4.15 (1H, sep, $J = 6.0$ Hz, $-\text{CH}<$), 3.65 (2H, s, $\text{CH}_2\text{-C=}$), 2.04 (2H, q, $J = 6.0$ Hz, CH_2), 1.06 (6H, d, $J = 6.0$ Hz, 2Me), 0.94 (3H, t, $J = 6.0$ Hz, Me); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 172.0 (C-2, C=O), 164.0 (C-4), 106.7 (C-3), 44.5 (C-5, CH_2), 41.0 (CH), 20.4 (2Me), 14.4 (CH_2), 13.1 (Me). FAB HRMS (acetone/NBA) calcd for $\text{C}_9\text{H}_{16}\text{NO}_2$ 170.1181 (M+H). Found 170.1171.

4-Hydroxy-1-(i-propyl)-3-propyl-3-pyrrolin-2-one (3p). $R_f = 0.55$ (EtOAc:hexane = 7:3 v/v); colorless needles (from EtOAc/hexane); mp 117–118 °C; IR (CHCl₃) ν 3300–3100 (OH), 1770, 1681 (C=O); ¹H-NMR (DMSO-*d*₆) δ 10.49 (1H, br s, OH), 4.16 (1H, sep, $J = 6.9$ Hz, -CH<), 3.66 (2H, s, CH₂-C=), 2.00 (2H, t, $J = 7.8$ Hz, CH₂), 1.36 (2H, m, CH₂), 1.07 (6H, d, $J = 6.9$ Hz, 2Me), 0.83 (3H, t, $J = 7.8$ Hz, Me); ¹³C-NMR (DMSO-*d*₆) δ 172.2 (C-2, C=O), 164.6 (C-4), 105.2 (C-3), 44.5 (C-5, CH₂), 41.1 (CH), 23.0, 21.2 (CH₂), 20.3 (2Me), 13.8 (Me). FAB HRMS (acetone/NBA) calcd for C₁₀H₁₈NO₂ 184.1338 (M+H). Found 184.1334.

1-Ethyl-4-hydroxy-3-methyl-3-pyrrolin-2-one (3q): $R_f = 0.32$ (EtOAc:hexane = 6:4 v/v); colorless microcrystals (from EtOAc/hexane); mp 121–122 °C; IR (CHCl₃) ν 3300–3100 (OH), 1772, 1685 (C=O); ¹H-NMR (DMSO-*d*₆) δ 10.54 (1H, br s, OH), 3.72 (2H, s, CH₂-C=), 3.27 (2H, q, $J = 7.2$ Hz, CH₂), 1.53 (3H, s, Me), 1.02 (3H, t, $J = 7.2$ Hz, Me); ¹³C-NMR (DMSO-*d*₆) δ 172.6 (C-2, C=O), 164.3 (C-4), 100.4 (C-3), 48.6 (C-5, CH₂), 35.4 (CH₂), 15.5, 6.2 (Me). FAB HRMS (acetone/NBA) calcd for C₇H₁₁NO₂Na 164.0687 (M+Na). Found 164.0692.

Ethyl 1-benzyl-4-hydroxy-3-pyrrolin-2-one-3-carboxylate (3r). $R_f = 0.68$ (CHCl₃:MeOH = 98:2 v/v); colorless microcrystals (from benzene/cyclohexane); mp 148–150 °C (lit., mp 148.5–150 °C); IR (CHCl₃) ν 3700–3200 (OH), 1706, 1602 (C=O); ¹H-NMR (CDCl₃) δ 7.36–7.23 (5H, m, arom H), 4.59 (2H, s, CH₂), 4.41 (2H, q, $J = 7.2$ Hz, CH₂), 3.85 (2H, s, CH₂-C=), 1.63 (1H, br s, OH), 1.41 (3H, t, $J = 7.2$ Hz, Me); ¹³C-NMR (CDCl₃) δ 183.8, 167.1 (C=O), 157.6 (C-4), 136.6 (arom C), 128.8, 128.1, 127.8 (arom CH), 98.6 (C-3), 61.3, 49.0, 45.4 (CH₂), 14.2 (Me).

1-Benzyl-4-hydroxy-3-phenyl-3-pyrrolin-2-one (3s). $R_f = 0.29$ (CHCl₃:MeOH = 98:2 v/v); colorless needles (from EtOH/hexane); mp 197–198 °C (lit., mp 197–198 °C); IR (KBr) ν 3500–2900 (OH), 1664 (C=O); ¹H-NMR (DMSO-*d*₆) δ 11.71 (1H, br s, OH), 8.02–7.14 (10H, m, arom H), 4.56 (2H, s, CH₂), 3.84 (2H, s, CH₂); ¹³C-NMR (DMSO-*d*₆) δ 170.9 (C-2, C=O), 166.9 (C-4), 138.0, 132.2, 128.5 (arom C), 127.6, 127.4, 127.0, 126.6, 125.5 (arom CH), 102.7 (C-3), 48.6 (C-5, CH₂), 44.4 (CH₂).

The physical data for the barbituric acids **5a–e** are given below.

1,3-Dimethyl-5-i-propylbarbituric acid (5a). Yield 90%; colorless microcrystals (from CH₂Cl₂/hexane); mp 105 °C; IR (CHCl₃) ν 1739, 1697, 1667 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 3.34 (1H, d, $J = 3.8$ Hz, -CH- of C-5), 3.29 (6H, s, 2 × CH₃), 2.60–2.54 (1H, m, CH₃-CH-CH₃), 1.06 (6H, d, $J = 6.8$ Hz, CH₃-CH-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.3, 151.7 (C=O), 55.2 (C-5), 36.8 (CH₃-CH-CH₃, *i*-Pr), 28.1 (CH₃), 19.4 (CH₃-CH-CH₃, *i*-Pr). Anal. Calcd for C₉H₁₄N₂O₃: C, 54.53; H, 7.12; N, 14.13. Found: C, 54.60; H, 7.06; N, 14.11.

5-Benzyl-1,3-dimethylbarbituric acid (5b). Colorless blocks (from CH₂Cl₂/hexane); mp 112 °C; IR (KBr) ν 1743, 1697, 1678 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.23–7.00 (5H, m, arom H), 3.78–3.75 (1H, m, -CH-), 3.44 (2H, d, $J = 4.6$ Hz, -CH₂-, Bn), 3.10 (6H, s, 2 × CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.3, 150.9 (C=O), 135.1 (arom C), 128.8, 128.5, 127.7 (arom CH), 50.6 (C-5), 37.8 (-CH₂-, Bn), 28.1 (CH₃). Anal. Calcd for C₁₃H₁₄N₂O₃•2/3H₂O: C, 60.46; H, 5.98; N, 10.85. Found: C, 60.54; H, 5.48; N, 12.22.

5-(4-Methoxyphenyl)methyl-1,3-dimethylbarbituric acid (5c). Colorless blocks (from CH₂Cl₂/hexane); mp 85 °C; IR (KBr) ν 1743, 1698, 1660 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 6.94–6.71 (4H, m, arom H), 3.73 (3H, s, CH₃O-), 3.71–3.67 (1H, m, -CH- of C-5), 3.38–3.36 (2H, m, -CH₂-), 3.10 (3H, s, CH₃), 3.04 (3H, s, CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.1, 158.7, 150.7 (C=O), 126.8 (arom C), 129.7, 113.6 (arom CH), 54.8 (CH₃O), 50.5 (C-5), 36.7 (-CH₂-), 27.8 (CH₃). Anal. Calcd for C₁₄H₁₆N₂O₄: C, 60.86; H, 5.84; N, 10.14. Found: C, 60.58; H, 5.43; N, 10.15.

5-(2-Methoxyphenyl)methyl-1,3-dimethylbarbituric acid (5d). Pale yellow needles (from CH₂Cl₂/hexane); mp 107 °C (decompd); IR (KBr) ν 1745, 1701, 1674 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.23–6.78 (4H, m, arom H), 3.80–3.77 (1H, m, -CH- of C-5), 3.73 (3H, s, CH₃O), 3.40 (2H, d, *J* = 5.4 Hz, -CH₂-), 3.10 (6H, s, 2 × CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.3, 157.1, 151.4 (C=O), 123.3 (arom C), 131.0, 123.3, 120.3, 110.0 (arom CH), , 55.0 (CH₃O), 50.0 (C-5), 33.3 (-CH₂-), 28.0 (CH₃). Anal. Calcd for C₁₄H₁₆N₂O₄: C, 60.86; H, 5.84; N, 10.14. Found: C, 60.86; H, 5.85; N, 10.07.

5-(2-Naphthyl)methyl-1,3-Dimethylbarbituric acid (5e). Yield 88%; colorless microcrystals (from CH₂Cl₂/hexane); mp 130 °C; IR (KBr) ν 1693, 1670 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.78–7.12 (7H, m, arom H), 3.84 (1H, t, *J* = 4.8 Hz, -CH- of C-5), 3.63 (2H, d, *J* = 4.8 Hz, -CH₂-), 3.09 (3H, s, CH₃), 3.01 (3H, s, CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.2 (C=O), 133.3, 132.7, 132.6 (arom C), 128.3, 127.9, 127.7, 127.5, 126.7, 126.4, 126.0 (arom CH), 50.7 (C-5), 37.5 (-CH₂-), 28.2 (CH₃).

The physical data for the quinolinone **7** are given below.

3-Butyl-4-hydroxy-2-quinolinone (7). Yellow microcrystals (from EtOAc/hexane); mp 196 °C (lit., mp 199–200 °C); IR (KBr) ν 3400–3000 (OH and NH), 1639 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.75 (1H, s, NH), 7.32–6.52 (4H, m, arom H), 2.01–1.97 (2H, m, -CH₂-C₃H₇), 0.86–0.73 (4H, m, -CH₂-CH₂-CH₂-CH₃), 0.30 (3H, t, *J* = 7.2 Hz, -C₃H₆-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 163.5 (C=O), 156.8 (C-4), 137.2, 115.3 (arom C), 129.4, 122.4, 120.7, 114.6 (arom CH), 111.7 (C-3), 30.3 (-CH₂-C₃H₇), 22.7 (-CH₂-CH₂-CH₂-CH₃), 22.2 (-CH₂-CH₂-CH₂-CH₃), 13.9 (-C₃H₆-CH₃).

The physical data for the malonamides **11a** and **11b** are given below.

2-Benzyl-N¹,N³-dipropylmalonamide (11a). 1.19 g (43% yield); *R*_f = 0.09 (CHCl₃); colorless microcrystals (from CHCl₃/hexane); mp 106–107 °C; IR (KBr) ν 3500–3200 (NH), 1630 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.28–7.16 (5H, m), 6.88 (2H, br. s), 3.24–3.08 (6H, m), 1.42 (4H sex, *J* = 7.2 Hz), 0.82 (6H, t, *J* = 7.2 Hz); ¹³C-NMR (75 MHz, CDCl₃) δ 170.4, 138.1, 128.8, 128.6, 126.8, 57.3, 41.3, 39.0, 22.5, 11.3.

2-Benzyl-N¹,N³-dibutylmalonamide (11b). 2.62 g (86% yield); *R*_f = 0.1 (CHCl₃); colorless needles (from CHCl₃/hexane); mp 108–109 °C; IR (KBr) ν 3400–3200 (NH), 1659 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.28–7.16 (5H, m), 6.81 (2H, br. s), 3.19–3.12 (6H, m), 1.37 (4H, quin, *J* = 7.2 Hz), 1.23 (4H, sex, *J* = 7.2 Hz), 0.87 (6H, t, *J* = 7.2 Hz); ¹³C-NMR (75 MHz, CDCl₃) δ 170.3, 138.1, 128.8, 128.5, 126.8, 57.2, 39.3, 39.0, 31.3, 19.9, 13.7. Anal. Calcd for C₁₈H₂₈N₂O₂: C, 71.02; H, 9.27; N, 9.20. Found: C, 70.84; H, 9.50; N, 9.26.

The physical data for the hydroperoxides **2a–h**, **4a–p**, **6a–e**, and **8** are given below.

4-Hydroperoxy-4-methyl-1,2-diphenylpyrazolidine-3,5-dione (2a). Colorless microcrystals (from CH₂Cl₂/hexane); mp 156–158 °C (decompd); IR (KBr) ν 3500–3200 (OOH), 1762, 1724 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 11.06 (1H, s, OOH), 7.34–7.16 (10H, m, arom H), 1.60 (3H, s, CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.9 (C=O), 134.9 (arom C), 129.1, 127.6, 123.4 (arom CH), 81.6 (>C-O), 17.7 (CH₃). Anal. Calcd for C₁₆H₁₄N₂O₄: C, 64.42; H, 4.73; N, 9.39. Found: C, 64.19; H, 4.70; N, 9.28.

4-Ethyl-4-hydroperoxy-1,2-diphenylpyrazolidine-3,5-dione (2b). Colorless microcrystals (from EtOAc/hexane); mp 141 °C; IR (KBr) ν 3600–3100 (OOH), 1762, 1724 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.67 (1H, s, OOH), 7.34–7.18 (10H, m, arom H), 2.04 (2H, q, J = 7.5 Hz, -CH₂-CH₃, Et), 1.03 (3H, t, J = 7.5 Hz, -CH₂-CH₃, Et); ¹³C-NMR (75 MHz, CDCl₃) δ 168.5 (C=O), 134.9 (arom C), 129.1, 127.4, 123.2 (arom CH), 85.8 (>C-O), 25.8 (-CH₂-CH₃, Et), 6.8 (-CH₂-CH₃, Et). Anal. Calcd for C₁₇H₁₆N₂O₄: C, 65.38; H, 5.16; N, 8.97. Found: C, 65.10; H, 5.13; N, 8.98.

4-Hydroperoxy-4-propyl-1,2-diphenylpyrazolidine-3,5-dione (2c). Colorless microcrystals (from Et₂O/hexane); mp 137 °C; IR (KBr) ν 3550–3200 (OH), 1764, 1726 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.95 (1H, s, OOH), 7.34–7.17 (10H, m, arom H), 1.98–1.93 (2H, m, -CH₂-C₂H₅), 1.47–1.41 (2H, m, -CH₂-CH₂-CH₃), 0.94 (3H, t, J = 7.2 Hz, -CH₂-CH₂-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.8 (C=O), 134.9 (arom C), 129.1, 127.5, 123.3 (arom CH), 85.3 (>C-O), 34.3 (-CH₂-CH₂-CH₃), 16.0 (-CH₂-CH₂-CH₃), 14.0 (-CH₂-CH₂-CH₃). Anal. Calcd. for C₁₈H₁₈N₂O₄: C, 66.25; H, 5.56; N, 8.54. Found: C, 65.99; H, 5.49; N, 8.45.

*4-Hydroperoxy-4-*i*-propyl-1,2-diphenylpyrazolidine-3,5-dione (2d)*. Colorless microcrystals (from Et₂O/hexane); mp 149–151 °C; IR (KBr) ν 3500–3150 (OOH), 1761, 1720 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.71 (1H, s, OOH), 7.37–7.16 (10H, m, arom H), 2.44 (1H, sep, J = 6.9 Hz, CH₃-CH-CH₃), 1.12 (6H, d, J = 6.9 Hz, CH₃-CH-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.6 (C=O), 135.0 (arom C), 129.0, 127.4, 123.1 (arom CH), 87.6 (>C-O), 33.4 (CH₃-CH-CH₃), 15.9 (CH₃-CH-CH₃). Anal. Calcd for C₁₈H₁₈N₂O₄: C, 66.25; H, 5.56; N, 8.58. Found: C, 65.97; H, 5.50; N, 8.49.

4-Butyl-4-hydroperoxy-1,2-diphenylpyrazolidine-3,5-dione (2e). Colorless microcrystals (from EtOAc/hexane); mp 124–126 °C (decompd), (lit, [7] mp 110–111 °C); IR (KBr) ν 3500–3150 (OOH), 1763, 1724 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.72 (1H, s, OOH), 7.34–7.18 (10H, m, arom H), 2.01–1.95 (2H, m, -CH₂-C₃H₇), 1.40–1.31 (4H, m, -CH₂-CH₂-CH₂-CH₃), 0.85 (3H, t, J = 6.9 Hz, -CH₂-CH₂-CH₂-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.7 (C=O), 134.9 (arom C), 129.0, 127.4, 123.2 (arom CH), 85.2 (>C-O), 32.0 (-CH₂-CH₂-CH₂-CH₃), 24.3 (-CH₂-CH₂-CH₂-CH₃), 22.5 (-CH₂-CH₂-CH₂-CH₃), 13.5 (-CH₂-CH₂-CH₂-CH₃). Anal. Calcd for C₁₉H₂₀N₂O₄: C, 67.05; H, 5.92; N, 8.23. Found: C, 66.83; H, 5.67; N, 8.05.

*4-*t*-Butyl-4-hydroperoxy-1,2-diphenylpyrazolidine-3,5-dione (2f)*. Colorless needles (from EtOAc/hexane); mp 173–178 °C (decompd); IR (KBr) ν 3500–3150 (OOH), 1747, 1717 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.51 (1H, s, OOH), 7.34–7.15 (10H, m, arom H), 1.19 (9H, s, 3 × CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.6 (C=O), 135.2 (arom C), 129.0, 127.2, 123.0 (arom CH), 89.1 (>C-O),

38.7 ((CH₃)₃C-), 24.3 ((CH₃)₃C-). Anal. Calcd for C₁₉H₂₀N₂O₄: C, 67.05; H, 5.92; N, 8.23. Found: C, 67.07; H, 5.86; N, 8.15.

4-Benzyl-4-hydroperoxy-1,2-diphenylpyrazolidine-3,5-dione (2g). Colorless blocks (from CH₂Cl₂/benzene); mp 103–107 °C (decompd); IR (KBr) ν 3400–3150 (OOH), 1753, 1705 (C=O); ¹H-NMR (300MHz, CDCl₃) δ 11.21 (1H, s, OOH), 7.24–6.80 (15H, m, arom H), 3.31 (2H, s, CH₂); ¹³C-NMR (75 MHz, CDCl₃) δ 167.7 (C=O), 134.0, 130.3 (arom C), 130.6, 128.8, 127.9, 127.6, 123.9 (arom CH), 86.7 (C-O), 38.0 (CH₂). Anal. Calcd for C₂₂H₁₈N₂O₄•2/3H₂O: C, 68.38; H, 4.69; N, 7.24. Found: C, 68.13; H, 4.55; N, 7.11. FAB HRMS (acetone/NBA) calcd for C₂₂H₁₉N₂O₄ 375.1345 (M+H). Found 375.1344.

X-ray crystallographic data of 2g (R = Bn). Empirical formula C₂₂H₁₈N₂O₄; formula weight 374.3894; colorless plates; crystal dimensions 0.25 × 0.50 × 0.10 mm; triclinic; space group *P*-1; *a* = 10.3233(5), *b* = 10.4273(4), *c* = 12.9985(6) Å, α = 95.792(1)°, β = 104.504(3)°, γ = 105.643(2)°, *V* = 1283.0(1) Å³, *Z* = 2; *D*_{calcd} = 1.272 g/cm³; *F*₀₀₀ = 518.00; μ (MoK α) = 0.85 cm⁻¹; $2\theta_{\max}$ = 55.0°; no. of reflections measured 11910; no. of observations (*I* > 3.00 σ (*I*), 2θ < 54.96°) 4605; no. of variables 442; reflection/parameter ratio 10.42; *R* = 0.038; *R*_w = 0.058; GOF 1.14. The crystallographic data (excluding structure factors) for the structure in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication number CCDC216767. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK [fax: +44(0)-1223-336033 or e-mail: deposit@ccdc.cam.ac.uk].

4-Cyclopentyl-4-hydroperoxy-1,2-diphenylpyrazolidine-3,5-dione (2h). Colorless microcrystals (Et₂O/hexane); mp 189 °C; IR (KBr) ν 3600–3200 (OOH), 1759, 1720 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.00 (1H, s, OOH), 7.35–7.16 (10H, m, arom H), 2.58–2.46 (1H, quin, *J* = 8.1 Hz, CH), 1.80–1.51 (8H, m, -(CH₂)₄); ¹³C-NMR (75 MHz, CDCl₃) δ 168.7 (C=O), 142.9, 135.1 (arom C), 129.1, 127.3, 123.1 (arom CH), 86.8 (>C-O), 43.5 (-(CH₂)₂-CH-(CH₂)₂-), 26.0, 24.8 (-(CH₂)₄-). Anal. Calcd for C₃₄H₃₂N₂O₄: C, 68.17; H, 5.72; N, 7.95. Found: C, 68.28; H, 5.72; N, 7.91.

1-Benzyl-3-hydroperoxy-3-methylpyrrolidine-2,4-dione (4a). Yield (221.1 mg, 94%); *R*_f = 0.67 (EtOAc:hexane = 8:2 v/v); colorless blocks (from Et₂O/hexane); mp 78–79 °C; IR (CHCl₃) ν 3400–3000 (OOH), 1786, 1666 (C=O); ¹H-NMR (CDCl₃) δ 11.28 (1H, s, OOH), 7.36–7.27 (5H, m, arom H), 4.79 (1H, d, *J* = 14.7 Hz, CH₂), 4.64 (1H, d, *J* = 14.7 Hz, CH₂), 3.75 (2H, s, CH₂), 1.39 (3H, s, Me); ¹³C-NMR (CDCl₃) δ 204.1 (C-4, C=O), 171.3 (C-2, C=O), 134.0, 129.1, 128.4, 128.3 (arom C), 82.5 (C-3), 53.4 (C-5, CH₂), 46.8 (PhCH₂), 16.7 (Me). FAB HRMS (acetone/NBA) calcd for C₁₂H₁₄NO₄ 236.0923 (M+H). Found 236.0935.

1-Benzyl-3-ethyl-3-hydroperoxy-pyrrolidine-2,4-dione (4b). Yield (236.8 mg, 95%); *R*_f = 0.53 (EtOAc:hexane = 6:4 v/v); colorless solid; mp 72–76 °C; IR (CHCl₃) ν 3500–3100 (OOH), 1782, 1693 (C=O); ¹H-NMR (CDCl₃) δ 11.45 (1H, s, OOH), 7.38–7.26 (5H, m, arom H), 4.73 (2H, s, CH₂), 3.74 (1H, d, *J* = 17.7 Hz, CH₂), 3.64 (1H, d, *J* = 17.7 Hz, CH₂), 1.85 (2H, q, *J* = 7.4 Hz, CH₂), 0.89 (3H, t, *J* = 7.4 Hz, Me); ¹³C-NMR (CDCl₃) δ 204.7 (C-4, C=O), 171.0 (C-2, C=O), 134.2, 129.1, 128.4, (arom C), 86.1

(C-3), 54.1 (C-5, CH₂), 46.8 (PhCH₂), 25.1 (CH₂), 16.70 (Me). FAB HRMS (acetone/NBA) calcd for C₁₃H₁₅NO₄Na 272.0899 (M+Na). Found 272.0906.

l-Benzyl-3-hydroperoxy-3-propylpyrrolidine-2,4-dione (**4c**). Yield (258.0 mg, 98%); *R_f* = 0.54 (EtOAc:hexane = 5:5 v/v); colorless solid; mp 112 °C; IR (CHCl₃) ν 3400–3100 (OOH), 1784, 1693 (C=O); ¹H-NMR (CDCl₃) δ 11.62 (1H, s, OOH) 7.37–7.27 (5H, m, arom H), 4.76 (1H, d, *J* = 14.7, CH₂) 4.67 (1H, d, *J* = 14.7 Hz, CH₂), 3.74 (1H, d, *J* = 17.7, CH₂), 3.64 (1H, d, *J* = 17.4 Hz, CH₂), 1.75 (2H, t, *J* = 8.4 Hz, CH₂), 1.31 (2H, m, CH₂), 0.88 (3H, t, *J* = 7.4 Hz, Me); ¹³C-NMR (CDCl₃) δ 204.8 (C-4, C=O), 171.1 (C-2, C=O), 134.2, 129.1, 128.4, 128.3 (arom C), 85.7 (C-3), 54.1 (C-5, CH₂), 46.8 (PhCH₂), 33.5, 15.9 (CH₂), 14.1 (Me). FAB HRMS (acetone/NBA) calcd for C₁₄H₁₇NO₄Na 286.1055 (M+Na). Found 286.1032.

l-Benzyl-3-butyl-3-hydroperoxypyrrrolidine-2,4-dione (**4d**). Yield (257.9 mg, 93%); *R_f* = 0.58 (EtOAc:hexane = 5:5 v/v); colorless solid; mp 80–81 °C; IR (CHCl₃) ν 3400–3100 (OOH), 1784, 1695 (C=O); ¹H-NMR (CDCl₃) δ 11.37 (1H, s, OOH), 7.38–7.26 (5H, m, arom H), 4.72 (2H, s, CH₂), 3.74 (1H, d, *J* = 17.6 Hz, CH₂), 3.64 (1H, d, *J* = 17.6 Hz, CH₂), 1.79 (2H, t, *J* = 8.1 Hz, CH₂), 1.18 (4H, m, 2CH₂), 0.84 (3H, t, *J* = 6.9 Hz, Me); ¹³C-NMR (CDCl₃) δ 204.8 (C-4, C=O), 171.0 (C-2, C=O), 134.2, 129.1, 128.4, 128.3 (arom C), 85.7 (C-3), 54.1 (C-5, CH₂), 46.8 (PhCH₂), 31.4, 24.4, 22.8 (CH₂), 13.6 (Me). FAB HRMS (acetone/NBA) calcd for C₁₅H₁₉NO₄Na 300.1212 (M+Na). Found 300.1219.

l-Benzyl-3-hexyl-3-hydroperoxypyrrrolidine-2,4-dione (**4e**). Yield (293.2 mg, 96%); *R_f* = 0.60 (EtOAc:hexane = 4: 6 v/v); colorless solid; mp 78–80 °C; IR (CHCl₃) ν 3400–3100 (OOH), 1784, 1691 (C=O); ¹H-NMR (CDCl₃) δ 11.56 (1H, s, OOH), 7.38–7.27 (5H, m, arom H), 4.75 (1H, d, *J* = 15.0 Hz, CH₂), 4.69 (1H, d, *J* = 15.0 Hz, CH₂), 3.75 (1H, d, *J* = 18.0 Hz, CH₂), 3.64 (1H, d, *J* = 18.0 Hz, CH₂), 1.78 (2H, t, *J* = 7.8 Hz, CH₂), 1.21 (8H, m, 4xCH₂), 0.85 (3H, t, *J* = 6.5 Hz, Me); ¹³C-NMR (CDCl₃) δ 204.8 (C-4, C=O), 171.1 (C-2, C=O), 134.2, 129.1, 128.4, 128.3 (arom C), 85.7 (C-3), 54.1 (C-5, CH₂), 46.8 (PhCH₂), 31.6, 31.3, 29.2, 22.4 (2C) (CH₂), 14.0 (Me). FAB HRMS (acetone/NBA) calcd for C₁₇H₂₃NO₄Na 328.1525 (M+Na). Found 328.1493.

l-Butyl-3-hydroperoxy-3-methylpyrrolidine-2,4-dione (**4f**). Yield (183.0 mg, 91%); *R_f* = 0.52 (EtOAc:hexane = 6:4 v/v); colorless liquid; IR (CHCl₃) ν 3400–3100 (OOH), 1784, 1693 (C=O); ¹H-NMR (CDCl₃) δ 11.75 (1H, s, OOH), 3.90 (2H, s, CH₂), 3.53 (2H, t, *J* = 6.9 Hz, CH₂), 1.60 (2H, m, CH₂), 1.36 (2H, m, CH₂), 1.35 (3H, s, CH₃), 0.96 (3H, t, *J* = 7.2 Hz, Me); ¹³C-NMR (CDCl₃) δ 204.9 (C-4, C=O), 171.3 (C-2, C=O), 82.2 (C-3), 53.9 (C-5, CH₂), 42.6, 28.6, 19.9 (CH₂), 16.6, 13.7 (Me). FAB HRMS (acetone/NBA) calcd for C₉H₁₅NO₄Na 224.0899 (M+Na). Found 224.0897.

l-Butyl-3-ethyl-3-hydroperoxypyrrrolidine-2,4-dione (**4g**). Yield (193.7 mg 90%); *R_f* = 0.66 (EtOAc:hexane = 6:4 v/v); colorless liquid; IR (CHCl₃) ν 3400–3100 (OOH), 1782, 1689 (C=O); ¹H-NMR (CDCl₃) δ 11.85 (1H, s, OOH), 3.90 (1H, d, *J* = 18.0 Hz, CH₂), 3.83 (1H, d, *J* = 18.0 Hz, CH₂), 3.63 (1H, m, CH₂), 3.48 (1H, m, CH₂), 1.79 (2H, q, *J* = 7.2 Hz, CH₂), 1.61 (2H, m, CH₂), 1.39 (2H, m, CH₂), 0.96 (3H, t, *J* = 7.2 Hz, Me), 0.88 (3H, t, *J* = 7.5 Hz, Me); ¹³C-NMR (CDCl₃) δ 205.3 (C-4, C=O), 170.7 (C-2, C=O), 85.7 (C-3), 54.5 (C-5, CH₂), 42.6, 28.4, 24.6, 19.7 (CH₂), 13.4, 6.7 (Me). FAB HRMS (acetone/NBA) calcd for C₁₀H₁₇NO₄Na 238.1055 (M+Na). Found 238.1036.

1,3-Dibutyl-3-hydroperoxy-pyrrolidine-2,4-dione (4h). Yield (231.1 mg, 95%); $R_f = 0.53$ (EtOAc:hexane = 5:5 v/v); colorless liquid; IR (CHCl₃) ν 3400–3100 (OOH), 1784, 1689 (C=O); ¹H-NMR (CDCl₃) δ 11.85 (1H, s, OOH), 3.88 (1H, d, $J = 18.0$ Hz, CH₂), 3.83 (1H, d, $J = 18.0$ Hz, CH₂), 3.61 (1H, m, CH₂), 3.48 (1H, m, CH₂), 1.73 (2H, t, $J = 7.8$ Hz, CH₂), 1.59 (2H, m, CH₂), 1.37 (2H, m, CH₂), 1.25 (4H, m, 2CH₂), 0.96 (3H, t, $J = 7.2$ Hz, Me), 0.85 (3H, t, $J = 6.8$ Hz, Me); ¹³C-NMR (CDCl₃) δ 205.3 (C-4, C=O), 170.8 (C-2, C=O), 85.3 (C-3), 54.4 (C-5, CH₂), 42.3, 30.9, 28.3, 24.2, 22.5, 19.6 (CH₂), 13.4, 13.3 (Me). FAB HRMS (acetone/NBA) calcd for C₁₂H₂₁NO₄Na 266.1368 (M+Na), Found 266.1375.

1-(t-Butyl)-3-hydroperoxy-3-methylpyrrolidine-2,4-dione (4i). Yield (181.0 mg, 90%); $R_f = 0.55$ (EtOAc:hexane = 7:3 v/v); colorless solid; mp 99–100 °C; IR (CHCl₃) ν 3400–3100 (OOH), 1782, 1697 (C=O); ¹H-NMR (CDCl₃) δ 11.27 (1H, s, OOH), 3.94 (2H, s, CH₂-C=O), 1.49 (9H, s, *t*-Bu), 1.32 (3H, s, Me); ¹³C-NMR (CDCl₃) δ 205.3 (C-4, C=O), 171.5 (C-2, C=O), 83.2 (C-3), 55.8 (>C<), 52.8 (C-5, CH₂), 28.1 (3CH₃), 16.9 (Me). FAB HRMS (acetone/NBA) calcd for C₉H₁₆NO₄ 202.1079 (M+H). Found 202.1069.

1-(t-Butyl)-3-ethyl-3-hydroperoxy-pyrrolidine-2,4-dione (4j). Yield (206.6 mg, 96%); $R_f = 0.64$ (EtOAc:hexane = 4:6 v/v); colorless solid; mp 130–131 °C; IR (CHCl₃) ν 3500–3200 (OOH), 1780, 1695 (C=O); ¹H-NMR (CDCl₃) δ 12.34 (1H, s, OOH), 4.08 (1H, d, $J = 18.3$ Hz, CH₂), 3.97 (1H, d, $J = 18.3$ Hz, CH₂), 1.58 (2H, q, $J = 7.5$ Hz, CH₂), 1.41 (9H, s, *t*-Bu), 0.77 (3H, t, $J = 7.5$ Hz, Me); ¹³C-NMR (CDCl₃) δ 207.7 (C-4, C=O), 169.2 (C-2, C=O), 85.3 (C-3), 54.5 (>C<), 53.3 (C-5, CH₂), 27.1 (3Me), 24.6 (CH₂), 16.9 (Me). FAB HRMS (acetone/NBA) calcd for C₁₀H₁₇NO₄Na 238.1055 (M+Na). Found 238.1045.

1-(t-Butyl)-3-hydroperoxy-3-propylpyrrolidine-2,4-dione (4k). Yield (227.1 mg, 99%); $R_f = 0.44$ (EtOAc:hexane = 4:6 v/v); colorless solid; mp 112–113 °C; IR (CHCl₃) ν 3450–3050 (OOH), 1782, 1685 (C=O); ¹H-NMR (CDCl₃) δ 11.59 (1H, s, OOH), 3.95 (1H, d, $J = 17.8$ Hz, CH₂), 3.84 (1H, d, $J = 17.8$ Hz, CH₂), 1.69 (2H, t, $J = 8.4$ Hz, CH₂), 1.49 (9H, s, *t*-Bu), 1.29 (2H, m, CH₂), 0.88 (3H, t, $J = 7.1$ Hz, Me); ¹³C-NMR (CDCl₃) δ 205.7 (C-4, C=O), 171.3 (C-2, C=O), 86.5 (C-3), 55.8 (>C<), 53.3 (C-5, CH₂), 33.7 (CH₂), 27.5 (3Me), 15.9 (CH₂), 14.1 (Me). FAB HRMS (acetone/NBA) calcd for C₁₁H₂₀NO₄ 230.1392 (M+H). Found 230.1042.

1-(t-Butyl)-3-butyl-3-hydroperoxy-pyrrolidine-2,4-dione (4l). Yield (240.9 mg, 99%); $R_f = 0.58$ (EtOAc:hexane = 4:6 v/v); colorless solid; mp 87–88 °C; IR (CHCl₃) ν 3400–3100 (OOH), 1782, 1685 (C=O); ¹H-NMR (CDCl₃) δ 11.47 (1H, s, OOH), 3.95 (H, d, $J = 18.0$ Hz, CH₂), 3.83 (1H, d, $J = 18.0$ Hz, CH₂), 1.72 (2H, t, $J = 7.2$ Hz, CH₂), 1.49 (9H, s, *t*-Bu), 1.25 (4H, m, 2CH₂), 0.82 (3H, t, $J = 7.1$ Hz, Me); ¹³C-NMR (CDCl₃) δ 205.8 (C-4, C=O), 171.2 (C-2, C=O), 86.4 (C-3), 55.8 (>C<), 53.6 (C-5, CH₂), 31.4 (CH₂), 27.5 (3Me), 24.4, 22.7 (CH₂), 13.5 (Me). FAB HRMS (acetone/NBA) calcd for C₁₂H₂₂NO₄ 244.1549 (M+H). Found 244.1521.

1-(t-Butyl)-3-hexyl-3-hydroperoxy-pyrrolidine-2,4-dione (4m). Yield (276.9 mg, 98%); $R_f = 0.45$ (EtOAc:hexane = 3:7 v/v); colorless solid; mp 94 °C; IR (CHCl₃) ν 3400–3100 (OOH), 1780, 1685 (C=O); ¹H-NMR (CDCl₃) δ 11.20 (1H, s, OOH), 3.94 (H, d, $J = 17.4$ Hz, CH₂), 3.83 (1H, d, $J = 17.4$ Hz,

CH₂), 1.71 (2H, m, CH₂), 1.49 (9H, s, *t*-Bu), 1.27 (8H, m, 4CH₂), 0.85 (3H, t, *J* = 6.9 Hz, Me); ¹³C-NMR (CDCl₃) δ 205.7 (C-4, C=O), 171.3 (C-2, C=O), 86.6 (C-3), 55.9 (>C<), 53.6 (C-5, CH₂), 31.8, 31.3, 29.2 (CH₂), 27.6 (3Me), 22.4, 22.3 (CH₂), 13.97 (Me). FAB HRMS (acetone/NBA) calcd for C₁₄H₂₆NO₄ 272.1862 (M+H). Found 272.1857.

3-Hydroperoxy-3-methyl-1-(i-propyl)pyrrolidine-2,4-dione (4n). Yield (177.8 mg, 95%); *R*_f = 0.41 (EtOAc:hexane = 7:3 v/v); colorless solid; mp 82–84 °C; IR (CHCl₃) ν 3400–3100 (OOH), 1784, 1685 (C=O); ¹H-NMR (CDCl₃) δ 11.65 (1H, s, OOH), 4.66 (1H, sep, *J* = 6.6 Hz, CH), 3.87 (1H, d, *J* = 17.7 Hz, CH₂), 3.80 (1H, d, *J* = 17.7 Hz, CH₂), 1.32 (3H, s, Me), 1.25 (3H, d, *J* = 6.6 Hz, Me), 1.22 (3H, d, *J* = 6.6 Hz, Me); ¹³C-NMR (CDCl₃) δ 204.9 (C-4, C=O), 170.5 (C-2, C=O), 82.6 (C-3), 49.0 (C-5, CH₂), 43.1 (CH), 19.43, 18.88, 16.36 (Me). FAB HRMS (acetone/NBA) calcd for C₈H₁₃NO₄Na 210.0742 (M+Na). Found 210.0740.

3-Ethyl-3-hydroperoxy-1-(i-propyl)pyrrolidine-2,4-dione (4o). Yield (195.2 mg, 97%); *R*_f = 0.50 (EtOAc:hexane = 7:3 v/v); colorless solid; mp 63–65 °C; IR (CHCl₃) ν 3600–3100 (OOH), 1782, 1685 (C=O); ¹H-NMR (CDCl₃) δ 11.66 (1H, s, OOH), 4.69 (1H, sep, *J* = 6.6 Hz, CH), 3.85 (1H, d, *J* = 17.4 Hz, CH₂), 3.71 (1H, d, *J* = 17.4 Hz, CH₂), 1.79 (2H, q, *J* = 7.3 Hz, CH₂), 1.26 (3H, d, *J* = 6.6 Hz, Me), 1.23 (3H, d, *J* = 6.6 Hz, Me), 0.78 (3H, t, *J* = 7.3 Hz, Me); ¹³C-NMR (CDCl₃) δ 205.3 (C-4, C=O), 170.1 (C-2, C=O), 86.4 (C-3), 49.7 (C-5, CH₂), 43.1 (CH), 24.7 (CH₂), 19.7, 18.9, 6.8 (Me). FAB HRMS (acetone/NBA) calcd for C₉H₁₅NO₄Na 224.0899 (M+Na). Found 224.0895.

3-Hydroperoxy-1-(i-propyl)-3-propylpyrrolidine-2,4-dione (4p). Yield (211.0 mg, 98%); *R*_f = 0.59 (EtOAc:hexane = 7:3 v/v); colorless solid; mp 78 °C; IR (CHCl₃) ν 3600–3100 (OOH), 1782, 1685 (C=O); ¹H-NMR (CDCl₃) δ 11.71 (1H, s, OOH), 4.68 (1H, sep, *J* = 6.6 Hz, CH), 3.85 (1H, d, *J* = 17.7 Hz, CH₂), 3.72 (1H, d, *J* = 17.7 Hz, CH₂), 1.71 (2H, t, *J* = 8.4 Hz, CH₂), 1.30 (2H, m, CH₂), 1.26 (3H, d, *J* = 6.6 Hz, Me), 1.22 (3H, d, *J* = 6.6 Hz, Me), 0.88 (3H, t, *J* = 7.2 Hz, Me); ¹³C-NMR (CDCl₃) δ 205.3 (C-4, C=O), 170.2 (C-2, C=O), 86.0 (C-3), 49.7 (C-5, CH₂), 43.0 (CH), 33.3 (CH₂), 19.6, 18.9 (Me), 15.9 (CH₂), 14.0 (Me). FAB HRMS (acetone/NBA) calcd for C₁₀H₁₇NO₄Na 238.1055 (M+Na). Found 238.1044.

5-Hydroperoxy-5-i-propyl-1,3-dimethylbarbituric acid (6a). Colorless oil; IR (CHCl₃) ν 3600–3100 (OOH), 1691 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.35 (1H, s, OOH), 3.37 (6H, s, 2 × CH₃), 2.38 (1H, sep, *J* = 6.9 Hz, CH₃-CH-CH₃), 0.97 (6H, d, *J* = 6.9 Hz, CH₃-CH-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.1, 150.7 (C=O), 89.5 (>C-O), 36.9 (CH₃-CH-CH₃), 29.0 (CH₃), 16.6 (CH₃-CH-CH₃). FAB HRMS (acetone/NBA) calcd for C₉H₁₄N₂O₅ 231.0903 (M+H). Found 231.0982.

5-Benzyl-5-hydroperoxy-1,3-dimethylbarbituric acid (6b). Colorless blocks (from CH₂Cl₂/hexane); mp 130 °C; IR (KBr) ν 3500–3200 (OOH), 1697, 1678 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.67 (1H, s, OOH), 7.28–7.26 (3H, m, arom H), 7.00–6.96 (2H, m, arom H), 3.25 (2H, s, CH₂), 3.13 (6H, s, 2 × CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.3, 149.4 (C=O), 130.2 (arom C), 129.1, 128.7, 128.6, 128.5 (arom CH), 87.2 (>C-O), 42.7 (CH₂), 28.6 (CH₃). Anal. Calcd for C₁₃H₁₄N₂O₅: C, 56.11; H, 5.07; N, 10.07. Found: C, 55.87; H, 5.01; N, 10.18.

5-Hydroperoxy-5-[(4-methoxyphenyl)methyl]-1,3-dimethylbarbituric acid (6c). Colorless blocks (from CH₂Cl₂/hexane); mp 117 °C; IR (KBr) ν 3600–3100 (OOH), 1728, 1692 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.85 (1H, s, OOH), 6.88–6.75 (4H, m, arom H), 3.76 (3H, s, CH₃O-), 3.19 (2H, s, CH₂), 3.16 (6H, s, 2 \times CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.3, 159.5, 149.5 (C=O), 121.9 (arom C), 130.2, 114.0 (arom CH), 87.2 (>C-O), 55.1 (CH₃O), 41.8 (-CH₂-), 28.6 (CH₃). Calcd for C₁₄H₁₆N₂O₆: C, 54.54; H, 5.23; N, 9.09. Found: C, 54.51; H, 5.21; N, 9.13.

5-Hydroperoxy-5-[(2-methoxyphenyl)methyl]-1,3-dimethylbarbituric acid (6d). Colorless plates (from CH₂Cl₂/hexane); mp 148–150 °C (decompd); IR (KBr) ν 3500–3200 (OOH), 1674 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 9.15 (1H, br, OOH), 7.27–6.78 (4H, m, arom H), 3.77 (3H, s, CH₃O), 3.27 (2H, s, -CH₂-), 3.14 (6H, s, 2 \times CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 167.9, 157.3, 150.0 (C=O), 131.7, 129.9, 120.5, 110.2 (arom CH), 119.1 (arom C), 86.9 (>C-O), 55.3 (CH₃O), 37.2 (-CH₂-), 28.6 (CH₃). Anal. Calcd for C₁₄H₁₆N₂O₆: C, 54.54; H, 5.23; N, 9.09. Found: C, 54.55; H, 5.15; N, 9.18.

5-Hydroperoxy-5-(2-naphthyl)methyl-1,3-dimethylbarbituric acid (6e). Colorless microcrystals (from CH₂Cl₂/hexane); mp 135 °C; IR (KBr) ν 3500–3200 (OOH), 1705, 1678 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 10.0 (1H, br, OOH), 7.80–7.03 (7H, m, arom H), 3.41 (2H, s, CH₂), 3.08 (6H, s, 2 \times CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 168.5, 149.4 (C=O), 133.2, 132.8, 127.6 (arom C), 128.7, 128.6, 127.8, 127.5, 126.9, 126.74, 126.4 (arom CH), 87.3 (>C-O), 42.9 (CH₂), 28.7 (CH₃). Anal. Calcd for C₁₇H₁₆N₂O₅: C, 62.19; H, 4.91; N, 8.53. Found: C, 62.45; H, 4.85; N, 8.54.

3-Butyl-3-hydroperoxy-1H-quinoline-2,4-dione (8). Pale yellow microcrystals (from EtOAc/hexane); mp 121 °C; IR (KBr) ν 3429 (OOH), 3350–3100 (NH), 1720, 1693, 1666 (C=O); ¹H-NMR (300 MHz, DMSO-*d*₆) δ 12.2 (1H, s, OOH), 11.0 (1, s, NH), 7.80–7.56 (2H, m, arom H), 7.14–7.03 (2H, m, arom H), 1.78–1.73 (2H, m, CH₂), 1.14–1.05 (4H, m, -CH₂-CH₂-CH₂-CH₃), 0.72 (3H, t, *J* = 6.9 Hz, -CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 193.9, 170.1 (C=O), 141.9, 119.0 (arom C), 136.8, 126.6, 122.8, 116.6 (arom CH), 88.6 (>C-O), 35.3 (-CH₂-CH₂-CH₂-CH₃), 24.2 (-CH₂-CH₂-CH₂-CH₃), 22.1 (-CH₂-CH₂-CH₂-CH₃), 13.4 (-CH₂-CH₂-CH₂-CH₃). FAB HRMS (acetone/NBA) calcd for C₁₃H₁₆NO₄ 250.1079 (M+H). Found 250.1062.

The physical data for the alcohols **9a–d**, **9g**, and **10b** are given below.

4-Hydroxy-4-methyl-1,2-diphenylpyrazolidine-3,5-dione (9a). Colorless microcrystals (from CH₂Cl₂/hexane); mp 172–173 °C; IR (KBr) ν 35100–3100 (OH), 1757, 1722 (C=O); ¹H-NMR (300MHz, CDCl₃) δ 7.33–7.17 (10H, m, arom H), 4.23 (1H, s, OH), 1.71 (3H, s, CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 170.8 (C=O), 135.1 (arom C), 129.0, 127.2, 122.7 (arom CH), 71.1 (>C-O), 23.4 (CH₃). Anal. Calcd for C₁₆H₁₄N₂O₃: C, 68.07; H, 5.00; N, 9.92. Found: C, 68.05; H, 4.97; N, 9.96.

4-Ethyl-4-Hydroxy-1,2-diphenylpyrazolidine-3,5-dione (9b). IR (KBr) ν 3500–3150 (OH), 1759, 1720 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.50–7.00 (10H, m, arom H), 4.54 (1H, s, OH), 2.05 (2H, q, *J* = 7.2 Hz, -CH₂-CH₃), 0.95 (3H, t, *J* = 7.2 Hz, -CH₂-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 170.6 (C=O), 134.7 (arom C), 128.9, 127.1, 122.8 (arom CH), 74.7 (>C-O), 30.6 (-CH₂-CH₃, Et), 7.0 (-CH₂-CH₃, Et).

4-Hydroxy-4-propyl-1,2-diphenylpyrazolidine-3,5-dione (9c). Colorless blocks (from Et₂O/hexane); mp 160 °C (lit, mp 167–169 °C); IR (KBr) ν 3500–3200 (OH), 1759, 1724 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.33–7.16 (10H, m, arom H), 3.73 (1H, s, OH), 2.06–2.00 (2H, m, -CH₂-C₂H₅), 1.48–1.40 (2H, m, -CH₂-CH₂-CH₃), 0.94 (3H, t, J = 7.2 Hz, -CH₂-CH₂-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 170.3 (C=O), 135.1 (arom C), 129.0, 127.2, 122.7 (arom CH), 74.0 (>C-O), 39.7 (-CH₂-CH₂-CH₃), 16.1 (-CH₂-CH₂-CH₃), 13.9 (-CH₂-CH₂-CH₃). Anal. Calcd for C₁₈H₁₈N₂O₃: C, 69.66; H, 5.85; N, 9.03. Found: C, 69.71; H, 5.81; N, 8.86.

4-Hydroxy-4-i-propyl-1,2-diphenylpyrazolidine-3,5-dione (9d). Pale yellow plates (from Et₂O/hexane); mp 175 °C (decompd); IR (KBr) ν 3550–3000 (OH), 1759, 1720 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.32–7.14 (10H, m, arom H), 3.85 (1H, br. S, OH), 2.34 (1H, sep, J = 6.9 Hz, CH₃-CH-CH₃), 1.15 (6H, d, J = 6.9 Hz, CH₃-CH-CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 17.1 (C=O), 134.8 (arom C), 129.0, 127.0, 122.7 (arom CH), 76.3 (>C-O), 36.2 (CH₃-CH-CH₃), 15.6 (CH₃-CH-CH₃). Anal. Calcd for C₁₈H₁₈N₂O₃: C, 69.66; H, 5.85; N, 9.03. Found: C, 69.91; H, 5.50; N, 9.01.

4-Benzyl-4-hydroxy-1,2-diphenylpyrazolidine-3,5-dione (9g). Colorless needles (from EtOAc/hexane); mp 200–210 °C (decompd) (lit, mp 206–208 °C); IR (KBr) ν 3500–3100 (OH), 1764, 1705 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.25–6.75 (15H, m, arom H), 4.88 (1H, s, OH), 3.44 (2H, s, CH₂); ¹³C-NMR (75 MHz, CDCl₃) δ 169.2 (C=O), 133.8, 131.6, 130.2 (arom C), 128.7, 128.6, 127.6, 127.2, 123.4 (arom CH), 75.8 (>C-O), 43.7 (CH₂). Anal. Calcd for C₂₂H₁₈N₂O₃: C, 73.73; H, 5.06; N, 7.82. Found: C, 73.47; H, 4.82; N, 7.76.

5-Benzyl-5-hydroxy-1,3-dimethylbarbituric acid (10b). Colorless blocks (from CH₂Cl₂/hexane); mp 112 °C; IR (KBr) ν 3500–3200 (OH), 1700, 1681 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.29–7.26 (3H, m, arom H), 6.98–6.95 (2H, m, arom H), 4.11 (1H, br, OH), 3.28 (2H, s, CH₂), 3.10 (6H, s, 2 × CH₃); ¹³C-NMR (75 MHz, CDCl₃) δ 169.9, 149.7 (C=O), 131.8 (arom C), 129.1, 128.5, 128.4 (arom CH), 76.6 (>C-O), 49.5 (CH₂), 28.4 (CH₃). Anal. Calcd for C₁₃H₁₄N₂O₄: C, 59.54; H, 5.38; N, 10.68. Found: C, 59.30; H, 5.33; N, 10.68.

The physical data for the alcohols **12a** and **12b** are given below.

2-Benzyl-2-hydroxy-N¹,N³-dipropylmalonamide (12a). Yield (18%); R_f = 0.79 (Et₂O); orange microcrystals; mp 74–76 °C; IR (KBr) ν 3398, 3360, 3312 (OH and NH), 1653 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.40 (2H, br. s), 7.27–7.23 (5H, m), 4.88 (1H, br. s), 3.21–3.15 (6H, m), 1.49 (4H, sex, J = 7.3 Hz), 0.85 (6H, t, J = 7.3 Hz); ¹³C-NMR (75 MHz, CDCl₃) δ 170.9, 134.9, 130.3, 128.0, 127.0, 77.8, 46.0, 41.5, 22.5, 11.3. Anal. Calcd for C₁₆H₂₄N₂O₃: C, 65.73; H, 8.27; N, 9.58. Found: C, 65.43; H, 8.56; N, 9.37.

2-Benzyl-N¹,N³-dibutyl-2-hydroxymalonamide (12b). Yield (25%); R_f = 0.79 (Et₂O); orange microcrystals; mp 75–77 °C; IR (KBr) ν 3377, 3314 (NH and OH), 1651 (C=O); ¹H-NMR (300 MHz, CDCl₃) δ 7.36 (2H, br.s), 7.26–7.22 (5H, m), 4.86 (1H, br. s), 3.23 (4H, q, J = 6.4 Hz), 3.14 (2H, s), 1.43 (2H, quin, J = 7.3 Hz), 1.27 (2H, sex, J = 7.3 Hz), 0.89 (3H, t, J = 7.3 Hz); ¹³C-NMR (75 MHz, CDCl₃) δ 170.9, 134.9, 130.3, 128.0, 127.0, 77.8, 46.0, 39.6, 31.3, 19.9, 13.7. Anal. Calcd for

$C_{18}H_{28}N_2O_3$: C, 67.47; H, 8.81; N, 8.74. Found: C, 67.40; H, 8.96; N, 8.73. FAB HRMS (acetone/NBA) calcd for $C_{18}H_{29}N_2O_3$ 321.2178 (M+H). Found 321.2184.

The physical data for the alcohols **14a** and **14b** are given below.

Methyl 3-hydroxy-2-oxotetrahydrofuran-3-carboxylate (14a). Yellow oil; IR (KBr) ν 3600–3100 (OH), 1768, 1720 (C=O); 1H -NMR (300 MHz, $CDCl_3$) δ 4.86 (1H, br, OH), 4.54–4.39 (2H, m, -CH₂-), 3.83–3.79 (1H, m, -CH₂-), 2.73–2.65 (1H, m, -CH₂-), 2.36 (3H, s, Me); ^{13}C -NMR (75 MHz, $CDCl_3$) δ 205.7, 174.8 (C=O), 81.2 (>C-O), 66.0 (-CH₂-), 33.8 (-CH₂-), 24.5 (CH₃).

Ethyl 2-hydroxycyclopentanone-2-carboxylate (14b). Pale yellow oil; IR (KBr) ν 3600–3100 (OH), 1757, 1732 (C=O); 1H -NMR (300 MHz, $CDCl_3$) δ 4.28–4.21 (2H, m, -CH₂-), 4.10 (1H, br, OH), 2.48–2.41 (3H, m, -CH₂-), 2.13–2.07 (3H, m, -CH₂-), 1.31–1.25 (3H, m, Me); ^{13}C -NMR (75 MHz, $CDCl_3$) δ 213.4, 171.5 (C=O), 79.7 (>C-O), 62.4 (-CH₂-), 35.7 (-CH₂-), 34.7 (-CH₂-), 18.2 (-CH₂-), 13.8 (CH₃).