

Table 1: Clinicopathological risk factors for survival in MF ICC patients

Factors		Number of patients	5-year survival rate (%)	Median survival time (months)	p-value
Overall		34	55.9	29	
Age (median: 65y.o.)	≥ 65	17	52.5	43	0.0119
	<65	17	39.0	17	
Sex	M	21	41.1	31	0.3746
	F	13	49.2	24	
CA19-9 (median: 53.8U/ml)	≥ 53.8	17	44.4	31	0.7175
	<53.8	16	35.1	28	
CEA (median: 1.6ng/ml)	≥ 1.6	14	23.5	16	0.0141
	<1.6	14	84.5	31	
Size (median: 45mm)	≥ 45	18	37.9	28	0.3642
	<45	16	54.3	33	
Intrahepatic metastses	Absent	21	51.2	36	0.1004
	Present	10	32.2	18	
Portal vein invasion	Absent	21	61.9	33	0.1676
	Present	13	46.2	20	
Bile duct invasion	Absent	14	64.3	31	0.2813
	Present	20	50.0	26	
Histological differentiation	well	8	58.3	41	0.3772
	mod/por	26	36.9	28	
Lymph node involvement	Absent	23	51.0	33	0.0284
	Present	11	24.0	10	
B or C hepatitis	Absent	23	41.9	29	0.9219
	Present	8	50.0	29	
Lymph node dissection	Absent	7	40.9	33	0.5233
	Present	26	46.4	28	
Sirius Red score (median: 25.4)	≥ 25.4	17	39.9	29	0.4619
	<25.4	17	46.8	29	
Platelet count (median 22.6X10 ⁴ /μl)	≥ 22.6	14	62.0	27	0.7604
	<22.6	13	60.0	31	